

Natural Occurrence Entomopathogenic Fungi: Beauveria Bassiana and Metharizium Anizopliae from the Population of Colorado Potato Beetle, Leptinotarsa Decemlineata (Coleoptera: Chrysomelidae) in Georgia

Nino Matcharashvili^{1*} and Medea Burjanadze^{2*}

¹PhD student of Faculty of Agricultural Sciences and Biosystem Engineering Georgian Technical University, Tbilisi, Georgia

²Professor, Faculty of Agricultural Sciences and Biosystem Engineering Georgian Technical University, Institute "Talga", Georgia

Corresponding author

Nino Matcharashvili and Medea Burjanadze, PhD student of Faculty of Agricultural Sciences and Biosystem Engineering Georgian Technical University, Tbilisi, Georgia and Professor, Faculty of Agricultural Sciences and Biosystem Engineering Georgian Technical University, Institute "Talga", Georgia.

Received: December 22, 2025; **Accepted:** December 30, 2025; **Published:** January 06, 2026

ABSTRACT

The paper demonstrated the Colorado potato beetle (CPB), *Leptinotarsa decemlineata* (Say) (Coleoptera: hrysomelidae) its natural enemies such as entomopathogenic fungi. Six isolates were obtained from larvae and adults of the population of *L. decemlineata*, at the different geographical and climatic zone of potato production. Morphological studies show that five isolates belong to the genus of Beauveria and one isolate of Metharhizium. Microscopic observation of colony growth and conidia arrangement showed general and typical characteristics of Beauveria bassiana in five isolates and one isolates Metharizium anizopliae.

Distribution of EPF among the insects, horizontal transmission, is a critical issue for the control of *decemlineata*. Experimental results show that adults of the *L. decemlineata* were infected with a fungal disease as a result of movement in the soil and spread naturally among them.

The beetles moved on the surface of the soil, as well as deep into the soil. On the 5th day few beetles appeared infected with mycosis; on the 9th day their number increased and infected beetles achieved around 50%; Finally, on the 14th day all the beetles were infected and their mortality reached 100%.

The identification and investigation of mycosis infection in the population of Colorado potato beetle, which has developed resistance to many insecticides used, is important to reduce economic losses and save the environment from pollution and is an important pathogen that can be used for the control of this insect.

Introduction

Potatoes are one of the most important staple foods in the world after wheat and rice and the third largest crop in the world in terms of human consumption [1]. In Georgia, the potato is the most widely grown crop in vegetable production [2]. A wide

range of insects can damage potatoes directly by feeding on the tubers and spoiling the crop or indirectly by feeding on leaves or stems. Tuber yield losses due to insects are estimated at 75% without crop protection [3].

Citation: Nino Matcharashvili, Medea Burjanadze. Natural Occurrence Entomopathogenic Fungi: Beauveria Bassiana and Metharizium Anizopliae from the Population of Colorado Potato Beetle, *Leptinotarsa Decemlineata* (Coleoptera: Chrysomelidae) in Georgia. *J Envi Sci Agri Res.* 2026. 4(1): 1-4.

DOI: doi.org/10.61440/JESAR.2026.v4.132

The Colorado potato beetle (CPB), *Leptinotarsa decemlineata* (Say) (Coleoptera)

hrysomelidae), is a serious pest of potato (*Solanum tuberosum* L.) in the USA and Europe, as well as in Asia [4,5]. The first considerable damage to the potato was reported in Colorado, USA [6]. By the end of the twentieth century, the pest had spread throughout Europe and Asia and continued to expand its geographic range to new regions of the world [7]. In Georgia, the pest was first discovered in 1965 in Gagra (Abkhazia) [8-10].

The beetle survives by feeding on cultivated and wild solanaceous plants, like potato, eggplant, and tomato. Both larvae and adults of CPB damage the leaves of potato plants. However, the main damage is caused by larval feeding [11].

At present, for the control CPB various chemical insecticides were used. The rapid development of resistance to various chemical insecticides requires the development of approaches to the control of *L. decemlineata* populations using natural enemies. The most widely used biological agents in biological control are entomopathogens [12]. Among them, entomopathogenic fungi (EPF) especially infects the pest through integument, their ease of production and high adaptability make them potentially more useful than the others. EPF has three mechanisms of action including causing mechanical damage to tissues and producing toxic metabolites. In this way, they can cause the direct death of insects or weaken them, and limit their vital activities [13].

The widely distributed entomopathogenic hyphomycete *Beauveria bassiana* (Balsamo) Vuillemin (Deuteromycota: Hyphomycetes) has been studied for microbial control of CPB [14-17]. *B. bassiana* shows efficient control against adult and all larval stages of CPB and provides an extremely high level of control during the potato growing season because it can continue to reproduce after application. However, its pathogenicity is affected by abiotic factors such as moisture, sunlight, and ultraviolet rays [18,19].

The objectives of this research were to find and identify an entomopathogenic fungus, spread in the population of *L. decemlineata*, at different geographical and different zones of potato production.

Material and Methods

Collection of Insects

Adults and larvae of *L. decemlineata* (I-V instars) were collected from four different geographical areas at different altitudes (600-1200 m a.s.l), potato fields: Kumisi, Tetrictkaro, Bulachauri, and Tianety, 2023-2024.

Isolation of Entomopathogenic Fungi

Single insects with the symptom's characteristic of entomopathogenic fungus have been transferred in the laboratory for the detection of fungi strains. Adults and larvae of CPB were placed in moister chambers to develop fungi mycelium. Fungi isolated from the insect cadaver were cultivated on artificial media and incubated at $23 \pm 2^\circ \text{C}$ for 12-15 days.

Colony Growth and Morphology

Colony descriptions and measurements were determined from

cultures grown on full-strength potato dextrose agar (PDA) (DifcoTM) on Petri dishes (ϕ 90 mm); The plates were incubated at 25°C in the dark at $23 \pm 2^\circ \text{C}$ in darkness from 10-14 day from inoculation. Terms and notations are used to describe colony coloration, hyphae, and conidia [20-22]. Microscopic measurements of conidiogenous cells and conidia were taken from PDA cultures at 5-15 days and images were acquired with a light microscope. Terminology for conidial shape follows hyphae and conidia measurements were taken [23].

Horizontal Transmission of *Beauveria Bassiana*

Experiments were conducted to determine the biological activity and distribution infection of the fungus *Beauveria bassiana* to the adult of the Colorado potato beetle. In the glass dishes (volume 2 liter), 1/3 volume with moister soil, were placed 30 adults of CPB, in each other container, with 3 replications.

Results and Discussion

Identification of Entomopathogenic Fungi

The samples were obtained from 4 different geographical sites at different altitudes (600-1200 meters above sea level (m a.s.l.) and climate zones, each representing a unique agricultural ecosystem in Georgia. In nature, we found infected larvae and adults (Figure 1) which were transferred to the laboratory for the study.



Figure 1: Occurrences infected larvae and Adults of *L. decemlineata* in nature: a - larvae with *Beauveria bassiana*; b – beetles with *Beauveria bassiana*; c – adult with *Metharhizium anisopliae*

Five isolates of *Beauveria bassiana* and one isolate of *Metharhizium anisopliae* were obtained from larvae and adults of *L. decemlineata*, as described in Table 1.

Table 1: Sites of investigation of *Leptinotarsa decemlineata* in cultivated potato field

Name of isolates	Name of sites	Geographical location (lat. N, long. E)	Altitude	Name of isolates	year
CPB 1	Kumisi	410 61' 36", 440 78' 21"	950	<i>B. bassiana</i>	2024
CPB 2	Tetrictkaro	410 55' 22", 440 48' 18"	1000	<i>B. bassiana</i>	2024
CPB 3	Shovi, GCM**	420 35' 36", 430 00' 88"	1800	<i>B. bassiana</i>	2023
CPB 4	Shovi, GCM	420 23' 76", 430 10' 13"	1650	<i>B. bassiana</i>	2023
CPB 5	Shovi, Glola, GCM	420 33' 36", 430 00' 58"	1730	<i>B. bassiana</i>	2023

CPB 6	Shovi, Glola, GCM	420 44' 07" , 430 77' 01"	1780	M. anizopliae	2023
-------	-------------------------	------------------------------	------	------------------	------

*LCM- Lesser Caucasian Mountain; GCM**- Grate Caucasian Range

Table 2: Morphological characterization of isolates Beauveria bassiana and Metarhizium anizopliae

Species	Colony description	Colony size (Ømm)	Size of conidia (µm)	Form of conidia
Beauveria bassiana (CPB 1, CPB 2, CPB 3, CPB 4, CPB,5)	cottony, powdery, velutinous, white or yellow white appressed to the agar surface	13–30	(1.5) 2.0 – 3.0 (4.0) x (1.5) 2.0 – 2.5 (- 3.0)	globose, subglobose, broadly ellipsoid forming short chains
Metarhizium anizopliae (CPB 6)	sub velutinous, velutinous to cottony, closely appressed to agar surface dark green	14–34	7.5-10.2 x 4.0–4.5 12.5- 15.0 x 2.5-3.0	elongated ellipsoidal to cylindrical

Colony of M.anizopliae growth on full-strength PDA 14–34 mm diam. at 12 days at 23oC.

Surface mycelium sub velutinous, closely appressed to agar surface, at the s yellowish green to dark green. Conidia aggregated as, 0.1 mm spherical clusters, white in mass, conidia occasionally. Forming a farinaceous surface layer in older cultures. The colony reverses a dark green color. The conidia, which were oval (n = 30), varied 9.0 (7.5–10.2) µm in length and 4.3 (4.0–4.5) µm in width. The matured spores are commonly elongated ellipsoidal to cylindrical, with dimensions 12.5-15.0 x 2.5-3.0 µm (Figure 2, c,d).




Figure 2: Isolates entomopathogenic fungi from L.decemlineata: a - colony of B.Bassiana; b - spores of B.Bassiana; c-colony of M.anizopliae ; d - spores of M.anizopliae.

Both fungi are known to produce cuticle-degrading enzymes that facilitate percutaneous infection of the host insect without the need for oral consumption by the target organism [24]. Fungi exhibit qualities that can make them ideal for this infection

strategy including a mild-acting nature, the ability to self-replicate, and the ability of fungal conidia to be spread by adults of L.decemlineata [25,26].



Figure 3: Horizontal transmission of mycosis caused by B.bassiana among adults of L.decemlineata: 5 days; b – 9 days; c – 14 days; d – infected insects in the soil.

Distribution of EPF among the insects is a critical issue for the control of L.decemlineata. Experimental results show that adults of the CPB were infected with a fungal disease as a result of movement in the soil (Figure 3). and spread naturally among them.

The beetles moved on the surface of the soil, also they were moving deep into the soil. On the 5th day of the beginning experiment, appeared single beetles infected with mycosis, On the 5th day of the experiment, single beetles infected with mycosis appeared, and on the 9th day their number increased and infected beetles achieved around 50%, Finally, on the 14th day all the beetles were infected and their mortality reached 100%.

Conclusion

The identification and investigation of mycosis infection in the population of Colorado potato beetle - L.decemlineata, which has developed resistance to many insecticides used, is especially important to reduce economic losses. It has been shown that Beauveria bassiana and Metarhizium anizopliae isolated from naturally infected insects is an important pathogen that can be used for the control of these insects. Subsequently, mycoses infection and its high efficiency were demonstrated in laboratory and pot experiments. In addition, the optimum condition for the occurrence of maximum pathogenicity of entomopathogenic fungus can be investigated.

References

1. FAOSTAT. Food and agriculture data. 2019.
2. GEOSTAT. 2020.
3. James C. Global Status of Commercialized Biotech/GM Crops; ISAAA: Ithaca, NY, USA. 2011.
4. Jolivet P. Le doryphore menace l'Asie Leptinotarsa decemlineata Say 1824 (Col. Chrysomelidae). Entomologist. 1991. 47: 29-48.
5. Weber D. Colorado beetle: pest on the move. Pestic Outlook. 2003. 14: 256-259.
6. Riley CV. Seventh Annual Report on the Noxious, Beneficial, and Other Insects of the State of Missouri, 1st ed.; Regan & Carter: Jefferson City, MO, USA. 1875.
7. Wang Y, Tang DX, Duan DE, Wang YB, Yu H. Morphology, molecular characterization, and virulence of Beauveria

pseudobassiana isolated from different hosts. *J Invertebr Pathol.* 2020. 172: 107333.

- 8. Burjanadze M. Microbiological method of controlling potato from Colorado Potato beetles in Eastern Georgia. Theses of PhD dissertation, Georgia. 1988. 146.
- 9. Goderdzishvili G, Parulava N. Colorado potato beetle in Georgia. *Georgian Agricultural.* 1981. 9: 32.
- 10. Antadze OK, Dmbadze AN, Parulava NI. Spread Colorado potato beetle in Georgia and its control. Proceeding of Georgia Research Scientific Institute Plants Protection. 1981. 31: 23-29.
- 11. Kivan M, Aysal T. Laboratuvar koşullarında *Perillus bioculatus* (F.) (Hemiptera: Pentatomidae)'un gelişme süresi ve ölüm oranı üzerinde ön çalışmalar. *Türkiye Biyolojik Mücadele Dergisi.* 2014. 5: 23-29.
- 12. Deacon JW. *Microbial Control of Plant Pests and Diseases.* Van Nostrand Reinhold Company, New York, US. 1983. 88.
- 13. Kulkarni SA. Biochemical and Molecular Studies of Chitin Deacetylase from *Metarhizium* species. Savitribai Phule Pune University, Biochemical Sciences Division CSIRNational Chemical Laboratory, (Unpublished) Ph.D Thesis, India. 2015. 196.
- 14. Lipa JJ. Progress in biological control of the Colorado beetle (*Leptinotarsa decemlineata*) in Eastern Europe. *Bull OEPP.* 1985. 15: 207-211.
- 15. Wraight SP, Ramos ME. Effects of inoculation method on the efficacy of wettable powder and oil dispersion formulations of *Beauveria bassiana* against Colorado potato beetle larvae under low humidity conditions. *Biocontrol Sci Technol.* 2017. 27: 348-363.
- 16. Ropék D, Kołodziejczyk M. Efficacy of selected insecticides and natural preparations against *Leptinotarsa decemlineata*. *Potato Res.* 2019. 62: 85-95.
- 17. Eski A, Bayramoglu Z, Sonmez E, Biryol S, Demir I. Evaluation of the effectiveness of entomopathogens for the control of Colorado potato beetle, *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). J Agr Sci Tech.
- 18. Fernandes ÉK, Rangel DE, Braga GU, Roberts DW. Tolerance of entomopathogenic fungi to ultraviolet radiation: a review on screening of strains and their formulation. *Curr Genet.* 2015. 61: 427-440.
- 19. Acheampong MA, Hill MP, Moore SD, Coombes CA. UV sensitivity of *Beauveria bassiana* and *Metarhizium anisopliae* isolates under investigation as potential biological control agents in South African citrus orchards. *Fungal Biol.* 2020. 124: 304-310.
- 20. Evlakhova AA. *Entomogenous Fungi. Classification, Biology, Practical Significance,* Nauka Press, Leningrad, Russia. 1974. 1-260.
- 21. Roberts, DW, Humber RA. Entomogenous fungi. In: G.T. Cole, B. Kendrick (Eds.), *Biology of Conidial Fungi*, Academic Press, New York, USA. 1981. 201-236.
- 22. Humber RA. Fungi Identification. In: *Manual of Techniques in Insect Pathology.* 1st ed. Chapter V-1, San Diego, CA, USA. 1997. 1: 153-185.
- 23. Vellinga EC. Glossary. In: C Bas, Th.W. Kuyper, M.E. Noordeloos, E.C. Vellinga (eds.), *Flora Agaricina Neerlandica, Critical monographs on families of agarics and boleti occurring in the Netherlands*, vol 1. A.A. Balkema, Rotterdam. 1988. 54-64.
- 24. Wright MS, Osbrink WLA, Lax AR. Transfer of entomopathogenic fungi among Formosan subterranean termites and subsequence mortality. *J Appl. Entomol.* 2002. 126: 20-23.
- 25. Grace JK, Zoberi MH. Experimental evidence for transmission of *Beauveria bassiana* by workers (Isoptera: Rhinotermitidae). *Sociobiology.* 1992. 20: 23-28.
- 26. Cheraghi A, Habibpour B, Mossadegh MS, Sharififard M. Horizontal Transmission of the Entomopathogen Fungus *Metarhizium anisopliae* in *Microcerotermes versus* Groups. *Insects.* 2012. 3: 709-718.