

# The Relationship between Cognitive Impairment in Psychiatric Patients and Readmission Rate to an Inpatient Facility

Cherilyn Isis Schuff and Patrick J Aragon\*

School of Psychology, Florida Institute of Technology, 150 W University Blvd, Melbourne, FL 32901, USA

## \*Corresponding author

Patrick J Aragon, School of Psychology, Florida Institute of Technology, 150 W University Blvd, Melbourne, FL 32901, USA.

Received: December 06, 2023; Accepted: December 19, 2023; Published: December 27, 2023

## ABSTRACT

The primary intention of this study was to further understand the impact of assessing cognitive impairment in psychiatric patients, as a mediating factor on readmission rates. Mild cognitive dysfunction impacts a patient's functional outcomes [1,2]. Little information exists to guide best practices in the treatment of adults with cognitive impairment who are hospitalized for acute conditions [2]. A cognitive impairment may impact patient prognosis and ability to function outside of a setting focused on stabilization. Neuropsychological testing is a valuable tool in predicting a patient's cognitive potential. However, creating a battery of tests that must be short enough to fit the needs of patients within an inpatient setting, allowing them to demonstrate their ability, while still providing enough information for accurate diagnosis and prognosis is a challenge.

Overall, this study aimed to investigate the utilization of brief cognitive screeners in identifying patients that may be more vulnerable to relapse and readmission. Cognitive impairment was assessed utilizing the Montreal Cognitive Assessment (MoCA). Readmission rate is defined by total admissions per individual patient. The use of community resources, such as medication management, case management, housing programs, and their relationship with readmission rates was examined. Archival data was taken from an inpatient facility's HIPAA-compliant electronic medical record (EMR) database. All personal identifying information was de-identified to ensure minimal risk of breaching confidentiality.

Although all objectives demonstrated significant correlations among the variables, there were no significant predictive models. This study found a positive linear relationship between admission rate and a higher MoCA score. This may partially be explained by Age ( $M = 69.09$ ,  $SD = 8.25$ ) having a negative relationship with inpatient readmission rate and a positive relationship with community resources. Within the inpatient setting, patients with higher cognitive functioning may be more aware of their mental health symptoms and are more likely to seek help. This may lead to more frequent admissions as they are able to seek help to address their mental health concerns. The relationship between patient demographics, community resource usage, and readmission rate was also examined.

Limitations of this study included the use of archival data, a small sample size, and a lack of diversity within the sample. Future directions for research include norming the MoCA for psychiatric populations and examining deficit patterns within the cognitive domains.

## Introduction

Inpatient mental health treatment is an essential, life-saving level of care. It aims to stabilize patients that are in crisis, experiencing acute psychiatric conditions with a relatively sudden onset, severe course, or a marked decompensation due to a more chronic condition [3]. Given the severity of a patient's presentation, inpatient mental health treatment within a psychiatric hospital is one of the most intensive levels of psychiatric care. These psychiatric facilities are licensed and operated as either state or public psychiatric hospitals, or state-licensed private psychiatric hospitals that primarily provide 24-hour inpatient care [4]. They also may also provide 24-hour residential care and/or

less than-24-hour care, which is known as outpatient, partial or day hospitalization [4]. Treatment is provided in a 24-hour secure locked, medically staffed hospital unit with a multimodal approach [4]. Routine psychiatric evaluations by a psychiatrist or nurse practitioner, twenty-four-hour psychiatric nursing care, medication evaluation, and a structured milieu in addition to an individual behavioral plan constitute treatment on the inpatient unit.

Typically, patients are admitted when they are an imminent danger to themselves or others, or they are unable to care for themselves. Sometimes they may be admitted voluntarily for

things such as medication adjustment, emotional distress, or related concerns. In either circumstance, it is determined that their psychiatric care or behavioral needs are unmanageable at any available lower levels of mental health care. Common reasons for inpatient psychiatric admission include suicide attempts or threats, drug overdoses, threats of violence, or psychosis [4]. Inpatient admission is intended to only be a few days to a few weeks. Specifically, the law states that a client is to be evaluated within 72 hours of admission, at which time the patient is either allowed to be released, signs in voluntarily, or is evaluated by a minimum of two treatment providers that deem the patient incompetent and unable to sign in voluntarily. To prevent any patient from being kept against their will, all patients with an involuntary status must be allowed to attend mental health court to ensure proper use of patient rights. Inpatient care is designed to be a short stop which removes a patient from a troublesome environment, de-escalating a crisis. This quick intervention is often a vital barrier to loss of life.

## Review of Literature

### *Admission Rates and Inpatient Demographics*

There are 1,756 24-hour psychiatric inpatient facilities within the United States [4]. According to the 2020 National Mental Health Services Survey (N-MHSS), there were only three states that each had 12 or more public psychiatric hospitals; New York, California, and Texas. Of all facilities surveyed by N-MHSS in 2020 that reported having an inpatient population, there were 85,948 total inpatient beds designated for mental health treatment services [4]. There were 12 states (California, Colorado, Connecticut, Florida, Hawaii, Iowa, Massachusetts, Minnesota, Montana, Nebraska, North Dakota, and Rhode Island) that had an inpatient bed utilization rate of 100% or more, meaning that there were more patients receiving mental health treatment services in inpatient settings than there were inpatient designated beds at any given time [4]. Of the 77,622 patients who received reported mental health treatment services in inpatient settings in April 2020, 27% were cared for in general hospitals, 38% were treated in public psychiatric hospitals, and 29% were admitted to private psychiatric hospitals [4]. Across all facility types, 58% of all patients who received mental health treatment services in inpatient settings in April 2020 were involuntarily admitted for care, 36% of patients were admitted with an involuntary non-forensic (non-criminal) legal status, and 22% were admitted with an involuntary forensic (criminal) legal status [4]. Public psychiatric hospitals reported the highest proportion of involuntarily admitted patients who received mental health treatment services in inpatient settings (89%) [4]. The lowest proportions of involuntary patients were reported by residential treatment centers for children (13%) and Veteran's Affairs Medical Centers (27%) [4].

Deinstitutionalization, or the movement that advocates for the transfer of psychiatric patients from long-term facilities back to their families or community-based alternatives, has led to changes in the utilization of inpatient units [5]. Shifts in demographic variables, disease-related, system-based, and economic factors have been observed in the treatment recipients of acute inpatient psychiatric units [4,5]. There has been an increase in admissions pertaining to patients with neurodevelopmental disorders, substance use disorders, and trauma and stressor-related disorders [4,5]. The most frequent diagnosis seen within the inpatient

population was depressive disorder (29.44 %), followed by schizophrenia spectrum and other psychotic disorders (17.12%), and substance-related and addictive disorders (14.76%) [4,5]. There was nearly a significant increase in personality disorders [5]. The use of restraints has also increased significantly [4,5]. Nearly a quarter of all patients were admitted more than one time [5]. Similarly, there has been an increase in patients being discharged to another facility [5].

In 2020, the mean age of patients admitted to inpatient psychiatric treatment was 40.33 years (median 40, range 18-91 years) [5]. The most common age range group of patients being admitted was between 36 and 55 years [5]. Compared to females, males were admitted at a marginally higher rate (51.97%) [5]. Over 80% of the inpatients were either single, widowed, or divorced [5]. Unemployment was present in 42.90% of the patients and 30.63% were disabled due to either a chronic mental illness, physical, or medical disability [5]. The most common aftercare plan is being discharged back to a patient's home or self-care and self-medication management [4,5]. The typical length of stay ranges from 1 to 136 days, with the median length being 5 days [5].

Patients are presenting to inpatient facilities with serious, chronic mental illness or disability, and are in severe psychiatric crisis. They may lack social, occupational, financial, and housing stability, which may obstruct patient progress and contribute to an increase in inpatient readmission rates [4,5].

### *Cognition*

Cognition, behavior, and individual potential are firmly interconnected in a cyclical relationship. Cognition influences behavior, and behavior, in turn, can either facilitate or hinder the realization of one's potential. Cognition can be defined as the advanced cortical functions that include thinking, remembering, planning, knowing, and analyzing [6]. Cognitive functioning encompasses attention, memory, psychomotor speed, executive functioning, language, and visuospatial ability. Intact cognition is necessary for an individual to be aware of their needs, goals, and to accomplish daily challenges [7]. Cognitive impairment contributes to the overall complexity of patient care and increases the patient's risk of poor functional outcomes during and after an acute episode of illness [1,2]. A mild cognitive impairment can impact an individual's ability to function, impede activities of daily living, and compromise the quality of a patient's life [9].

Mild cognitive dysfunction increases the risk for psychiatric inpatient hospital readmissions as it impacts a patient's functional outcomes [1,2]. A matched case-control study looked at patients, 65 years or older, who were readmitted to the hospital within 30 days of discharge in a Medicare-managed care plan. Results indicated that these patients consumed additional health care resources, such as emergency visits to the medical hospital, and may be appropriate targets for interventions designed to reduce readmission. Little information exists to guide best practices in the treatment of older adults with cognitive impairment who are hospitalized for acute conditions [2]. If obtained, level of cognitive functioning can potentially inform intervention, predict functional outcome and readmission rates [2]. Improved recognition of cognitive impairment within an inpatient setting may decrease the number of inpatient readmissions. Cognitive

impairment impacts a patient's self-care; impairment in self-care significantly predicted readmission within a 30-day period [9].

### ***Cognitive Functioning in Psychotic Disorders***

Cognitive functioning in psychotic disorders, such as schizophrenia, schizoaffective disorder, and other related conditions, is often impaired and can have a significant impact on an individual's daily life. Cognitive dysfunction is a core feature of schizophrenia that is relevant to an increasing portion of the inpatient population (17.21%) [1,5]. Cognitive impairments are reported in approximately 75% of individuals diagnosed with schizophrenia [10]. Deficits have been observed to be moderate to severe across several domains, including attention, working memory, verbal learning and memory, and executive functions [1]. Cognitive deficits pre-date the onset of active psychosis and are stable throughout the course of the illness in most patients [1]. Over the past decade, the focus on these deficits has increased with the recognition that they are consistently the best predictor of functional outcomes across activities of daily living, treatment compliance, and medication adherence [1]. In addition to medication management, providers question the possibility for further intervention or compensatory strategies with cognitive re-training.

There is some debate as to whether cognitive impairments in schizophrenia are generalized or impact specific domains and can be recognized and profiled via neuropsychological assessment [1]. This argument is confounded by the fact that there is not a neuropsychological pattern of schizophrenia in testing as there sometimes is with other disorders, such as Alzheimer's Disease [11]. While there may be domain-specific deficits in patients presenting with psychotic disorders, the degree and pattern of impairment can vary from person to person [1,11]. Schizophrenia is a heterogeneous disorder, meaning that it manifests differently in different individuals. Some people with schizophrenia may have more pronounced cognitive deficits, while others may have relatively intact cognitive functioning. The heterogeneity suggests that cognitive profiles can vary widely among individuals with schizophrenia, which has implications for treatment planning and rehabilitation [11].

Some evidence has suggested that there are discrete domains of cognitive impairment within patients that have schizophrenia [11]. There are mild to moderate deficits in attention, verbal fluency, working memory, and processing speed, with superimposed severe deficits in declarative verbal memory and executive functioning [11]. Other work has suggested that discrete cognitive domains have differential correlates with symptom and functional domains [11]. In other words, patients with schizophrenia demonstrate at least some cognitive impairment, but, like other aspects of the illness, the severity and presentation of these impairments vary across patients [11]. A unique feature of cognitive deficits, as compared to other characteristics of schizophrenia, is that they remain relatively stable within the same patient over time [11]. The presentation of cognitive deficits of patients with schizophrenia are generally consistent in severity and topography across changes in a patient's clinical status [1,11].

## **Cognitive Deficits by Domain**

### ***Attention***

Impaired attention is considered a primary cognitive deficit in schizophrenia [1,11]. Individuals who are genetically predisposed to schizophrenia have a poor ability to maintain their attention even prior to their first psychotic episode [11,12]. Attentional impairments are typically present and are of moderate severity by the first onset of psychosis [1,13]. They include deficits in sustained attention, selective attention, and divided attention. Sustained attention, or the ability to maintain focus on a task or activity for an extended period, is often impaired in individuals with psychotic disorders. They may have difficulty concentrating on a single task, which can lead to problems with productivity, completing activities of daily living, or treatment adherence [1,13].

Selective attention refers to the ability to focus on specific stimuli while ignoring irrelevant information. People with psychotic disorders may have difficulty filtering out irrelevant sensory input, leading to sensory overload and increased distractibility [1,13].

Divided attention involves simultaneously processing multiple tasks or stimuli. Individuals with psychotic disorders may struggle with divided attention, finding it challenging to manage multiple demands or switch between tasks effectively [1,13].

Auditory hallucinations, a common symptom in schizophrenia, can interfere with sustained auditory attention. Individuals may be preoccupied with hearing voices, making it difficult to concentrate on other auditory stimuli or tasks [1,13].

Overall, attentional deficits can contribute to the severity of positive symptoms in psychotic disorders, such as hallucinations and delusions. For example, individuals may be more prone to misinterpret sensory information or be less able to discern between reality and hallucinatory experiences. Negative symptoms, which involve diminished emotional expression, social withdrawal, and reduced motivation, can also be influenced by attentional deficits. Impaired attention may limit an individual's ability to engage in social interactions and perform daily activities [1,13].

### ***Verbal Fluency***

Patients with schizophrenia have difficulties producing speech on demand [1]. Verbal fluency tests assess their ability to produce words from a specific phonological or semantic category, such as the Controlled Oral Word Association Test [1]. These tests reveal both poor encoding and storage of verbal information in addition to inefficient retrieval of information from semantic networks in patients with schizophrenia [1,14-16]. Individuals with psychotic disorders often produce fewer words than healthy controls on verbal fluency tasks. They may struggle to generate enough words within the time constraints. Verbal fluency deficits may be related to difficulties in accessing and retrieving words from their mental lexicon. This can lead to pauses, hesitations, and difficulty finding appropriate words. In category-based tasks, individuals may have difficulty shifting between different categories (e.g., switching from animals to fruits). In letter-based tasks, they may struggle to switch to words that begin with a new letter [14]. Successful performance on verbal fluency tasks often involves clustering words by related concepts (e.g., naming

several types of fruits in succession) and efficient switching between categories or letters. Individuals with psychotic disorders may exhibit deficits in these strategies [14]. Deficits in verbal fluency are also associated with poor interpersonal functioning [17].

### **Executive Functioning**

Executive functioning encompasses a wide range of cognitive processes that ultimately result in purposeful, goal-directed behavior [1]. Executive functions enable a patient to plan, follow multiple-step directions, in addition to displaying self-control. Executive functioning deficits can impact various aspects of daily life, including the ability to plan and carry out tasks, make sound decisions, and adapt to changing circumstances.

Studies using formal neuropsychological assessments have found that many patients with schizophrenia have difficulties with most or all these component processes [1]. For example, patients have a difficult time forming a conceptual framework to understand ambiguous stimuli due to inflexible, rigid thinking [1]. If a concept is comprehended, patients with schizophrenia have trouble adapting to changes in the environment that require different behavioral responses [1]. Rigid thinking is highly correlated with occupational difficulties [1]. Planning is an additional component of executive functioning that is often found to be impaired in patients with schizophrenia [1,16]. Self-care, social, interpersonal, community, and occupational functions are all associated with executive functioning in schizophrenia [1,18]. Importantly, executive functions are also associated with treatment success [1]. Impairments in this domain are associated with less engagement in therapy medication compliance [19] and longer hospital stays.

Recognizing specific cognitive deficits is important because it can inform targeted cognitive remediation interventions. Cognitive remediation programs aim to improve cognitive functioning in specific domains, helping individuals with schizophrenia develop strategies to compensate for their deficits. These programs often involve cognitive training exercises and psychoeducation for both the patient and their caretakers to enhance cognitive skills.

Executive functions encompass a range of cognitive processes, including planning, organizing, initiating, and inhibiting actions, shifting between tasks, problem-solving, working memory, and cognitive flexibility. In individuals with psychotic disorders, deficits in executive functioning are often characterized by difficulties in one or more of these areas.

These deficits can hinder independent living, educational attainment, and occupational functioning. Executive functioning deficits have been linked to both positive symptoms (e.g., hallucinations and delusions) and negative symptoms (e.g., diminished emotional expression and social withdrawal) in psychotic disorders. Impairments in executive functions may contribute to difficulties in interpreting and responding to reality, as well as limitations in social interactions and goal-directed activities.

### **Patient Readmission to Inpatient Psychiatric Hospitals**

#### **Readmission Rates**

It is estimated that about one-third of patients admitted to inpatient psychiatric services will probably be readmitted within

a year [20]. The readmission rate may be used to measure the extent of the quality of care rendered to patients which is also the focus of interest for all health sector policymakers [21]. The readmission rate is also used as the determinant of quality of care for measuring outcomes such as continuity of care and follow-up services.

The concept and definition of the readmission period varies between studies. For example, across studies it has been defined as readmission within 90 days, readmission within 3 years, admission three or more times within 30 months, readmission within 6 months, and admittance of 3 or more times within 2.5 years. This has resulted in a variance in the reported rates of readmission in different studies as 14%, 16%, 20-30%, and 45-53%, respectively.

#### **Factors in Readmission Rates**

Poor treatment adherence, involuntary first admission, substance, alcohol, and drug abuse all have been identified as related factors to the readmission of patients to inpatient psychiatric facilities. Other demographic characteristics such as marital status, unemployment, retirement, and gender, are also considered relevant factors that can influence readmission. Specific diagnoses may be correlated with a higher readmission rate. For example, the diagnosis of schizophrenia has been reported as a significant factor in readmission. The diagnoses of bipolar disorder, psychotic disorders, and mood disorders also have been indirectly connected with readmission. Diagnosis does mean that readmission is inevitable. However, as discussed previously, some diagnoses are correlated to deficits across cognitive domains creating an obstacle for treatment adherence. The severity of symptom presentation may make it more likely that a patient is readmitted to an inpatient facility [1,18]. In some studies, diagnoses with more psychiatric comorbidities were also a factor associated with readmission. Suicidal ideation or thoughts of self-harm, history of psychological problems in childhood, and the first episode of the psychiatric disorder under 18 years have all been revealed as related factors for readmission. The length of stay in the inpatient hospital has also been observed to impact readmission. The briefer the psychiatric inpatient admission, the more that it is associated with a higher likelihood of readmission. This can affect the quality of care and increase the costs of service to the provider. Identifying and understanding the influential factors related to psychiatric readmission rates may help to implement and manage preventative interventions. Identifying predictors of readmission at the inpatient level of care can inform health policies and quality improvement interventions to mitigate cost and the burden to systems and families.

#### **Assessment of Cognitive Impairment**

Best practice for thoroughly evaluating cognitive impairment includes the administration of a tailored neuropsychological battery. Neuropsychological test batteries typically focus on the assessment of multiple cognitive domains which may include learning, memory, language, processing speed, executive functioning, attention, and visuospatial ability. The main advantage of a full neuropsychological battery is that they may identify patterns of strengths and weaknesses across multiple functional domains which aids diagnosis and better predicts prognosis. They may also be able to demonstrate a patient's ability to care for



themselves. However, full neuropsychological batteries are rarely assessed formally in acute care, or psychiatric inpatient settings. This is primarily because they can be extensive, time consuming, difficult to administer with a patient in an acute state, may require specialized training for interpretation, and are unavailable in some facilities. Stabilization is understandably the focus for patients in these settings. Therefore, it is difficult to track the cognitive change during and between admissions within patients in these settings [2]. The negative symptoms of a psychotic disorder can be difficult to discern from the deficits of cognitive decline, however, can drastically impact treatment planning and success. A cognitive impairment may impact patient prognosis and ability to function outside of a setting focused on stabilization. Therefore, neuropsychological testing is required to thoroughly answer this question. However, creating a battery of tests that must be short enough to fit the needs of the patient and allow them to demonstrate their ability, while still providing enough information for accurate diagnosis and prognosis is a challenge.

### Assessments Utilized in Psychiatric Settings

The global cognitive deficit, or deficits across cognitive domains, of patients with schizophrenia is an average of 1 to 2 standard deviations (SD) below the healthy control mean. As previously reviewed, patients with schizophrenia are more likely to be impaired in areas of verbal skills, memory, attention, processing speed, and executive functions, with deficits up to 2.5 SD below control subjects [22,23]. These neurocognitive deficits could affect an individual's potential across different functional domains, such as occupational and interpersonal functioning, the completion of activities of daily living, and independent living [14]. Classic features of psychotic disorders that may be present are delusions, hallucinations, disturbed thinking processes, flattening of affect, and abnormal behaviors. Symptoms can be either persistent or episodic.

### Brief Cognitive Batteries

Brief cognitive batteries, such as the Brief Assessment of Cognition in Schizophrenia (BACS), the Brief Cognitive Assessment (BCA), and the Brief Cognitive Assessment Tool for Schizophrenia (B-CATS) have been developed to assess cognitive functioning in individuals with schizophrenia. Unfortunately, their psychometric properties are not well established, and they are not typically used in clinical practice. The disadvantage to using the BACS is that it requires approximately 35 minutes of administration time with the patient and additional scoring time for the clinician. The attention span of patients in an inpatient psychiatric setting can be variable, and therefore, a short battery is advantageous to assessment [25]. Although they only require a 12-to-15-minute administration time, the BCA and B-CAT present some scoring limitations for the clinician [25]. Raw scores are based on cognitive domains and need to be converted into z-scores. There is no direct total score or cut-off score that is available for an easy assessment of the cognitive functioning, limiting the test interpretation [25].

### The Mini-Mental State Examination (MMSE)

The Mini-Mental State Examination (MMSE), a widely utilized, first-choice screening tool across professions, has been used in clinical practice to detect cognitive impairment in relation to neurological disorders. While previous studies support the association of MMSE with age and education, few studies have

examined aspects like the association of cognitive impairment with clinical factors such as age of illness onset, types of antipsychotic medications [26], and duration of illness. Several studies have also shown that the MMSE is largely insensitive to identifying mild cognitive impairment and that it may not be a useful tool in the presence of mood disorders [27,28]. In addition, the MMSE was found to have a poor sensitivity for detecting cognitive disturbances in community mental health centers [29].

### The Texas Functional Living Assessment (TFLS)

The Texas Functional Living Scale (TFLS) is a performance-based measure of functional abilities with an emphasis on instrumental activities of daily living skills that is brief and weighted toward cognitive tasks [30]. There are 24 ordinary activities to complete, such as using a telephone, reading a calendar, telling time, calculating money, and memory tasks. The TFLS demonstrated the highest predictive ability for examining activities of daily living, however it may be too long to administer within an inpatient psychiatric facility [30].

### The Montreal Cognitive Assessment (MoCA)

The Montreal Cognitive Assessment (MoCA) has been utilized with individuals in acute inpatient psychiatric settings as a screening instrument with variable results [31]. It was developed in 2005 for the purpose of detecting mild cognitive impairment (MCI) and its' results have been shown to be highly sensitive and specific in the older adult population [32]. The MoCA is a brief neuropsychological screener that is simple enough to be administered at a patient's bedside. Total administration time typically runs about 10 to 15 minutes. The MoCA assesses short-term memory, attention, working memory, and executive functions, which are commonly affected in patients admitted to a crisis stabilization unit, or inpatient psychiatric hospitals. Scores on the MoCA range from zero to 30 points, with a score of 25 or lower indicating a cognitive impairment [31]. This cut-off score of 25 or lower is widely recognized as a threshold for detecting cognitive impairment and potential dementia. Practice effects have been minimized as there are three versions of the MoCA, which test the same domains but differ in content [31]. The alternative versions of the MoCA present comparable reliability to the original test [33]. The MoCA has also been translated into more than 60 languages [31]. In addition, training is available on the MoCA website. Certification for the MoCA was made a mandatory requirement as of September 2019 [31]. The process of training is standardized so that testing is consistent among multiple raters, minimizing errors, and thus minimizing liability and misdiagnosis [31].

### A Review of MoCA Literature

The literature on the MoCA as a diagnostic screener for inpatient psychiatric settings is mixed. The MoCA was utilized in diagnosing cognitive impairment in psychiatric inpatients [34]. A threshold score of 23, which is not the typical cut score, represented good sensitivity and specificity (0.82 and 0.70) in diagnosing neurocognitive disorders within the inpatient population [34]. The data has suggested that the total score on the MoCA is the most accurate in distinguishing neurocognitive disorders, such as dementia, from anxiety [34]. A MoCA threshold score of 23 was the least accurate in terms of differentiating neurocognitive disorders from psychotic disorders [34]. 77 patients diagnosed with schizophrenia were given the MoCA,

25.9% scores on MoCA between 0 and 19, 46.7% presented a score between 20 and 26, and 27.2% presented a score of 27 or more representing large range of scores [34].

The MoCA was utilized to detect cognitive impairment in 140 patients with long-term psychotic disorders [35]. Cognitive abilities were assessed with the MoCA test and the Matrices Consensus Cognitive Battery [35]. The Cronbach's alpha of MoCA was 0.76, suggesting only moderate reliability [35]. Regarding the cognitive domains assessed, the MoCA's executive/visuospatial subtest showed significant associations with all the domains except social cognition and attention/vigilance [35]. There have been more cases of Mild Cognitive Impairment that were detected when using the MoCA than when using the MMSE. As a cognitive assessment screening instrument, the MoCA showed adequate concurrent validity when compared with the scores obtained with the MMSE and with the global severity observed in the Positive and Negative Syndrome Scale (PANSS) assessment.

The correlation between the MoCA total score and the length of stay in an inpatient psychiatric facility was negative [25]. A higher MoCA score was associated with a shorter length of stay [25]. The average MoCA total score was 20.26 (SD = 5.63), ranging from 3 to 29 [25]. About 85% of the scores fell below the suggested 26 cut-off score, indicating most of the patients had some degree of cognitive impairment [25]. On average, patients with schizophrenia performed worse on every single item as well as the total score compared to existing MoCA normative data of normal controls [32]. The patients performed the worst on the delayed recall task with most patients (92%) unable to score full points [25]. Other items that presented difficulty included abstraction, orientation, sentence repetition, serial 7 subtractions, verbal fluency, and trail making [25]. Most of the patients (84%) correctly performed the naming task [25].

When validating MoCA against the MCCB, five out of seven MoCA subtests (visuospatial-executive, attention, language, abstraction, and delayed recall) were significantly associated with the MCCB subtests [35]. The naming and orientation MoCA subtests were not correlated with the MCCB subtests [35]. In addition, the social cognition and vigilance domains assessed by MCCB were the less correlated with MoCA because it does not assess social cognition or the ability of the patient to maintain sustained attention [35]. Moreover, the MoCA does not specifically assess processing speed, which is one of the core cognitive impairments in patients with schizophrenia [35].

When compared to other short batteries such as MMSE, investigations of the MoCA in psychotic spectrum disorders have proved its superiority. The MoCA contains specific subtests addressing executive functions and attention.

A possible explanation for the different results obtained across studies in patients with schizophrenia could be that MoCA does not evaluate some specific cognitive deficits of schizophrenia [35]. Despite this, the items of MoCA could be useful in older patients, which can present multiple comorbidities such as increased vulnerability to Alzheimer's disease, cardiovascular risk factors, and cerebrovascular disease [35]. Cognitive impairment is associated with illness severity and negative symptoms [25]. There

was no significant decline or improvement in cognitive impairment during hospital stay in patients with severe schizophrenia, even though their illness severity and clinical symptoms improved significantly at discharge [25]. Cognitive impairment is relatively stable over different test-retest intervals and across psychotic state changes [25]. However, it is worth noting that orientation scores improved from admission to discharge [25].

Although the MoCA has the potential to be a promising screening test for patients with schizophrenia, when used at the recommended cutoff score of 26, it may not be the best discriminating tool for this specific population [25,35]. While clinicians may prefer a test with high sensitivity, this could increase the number of patients referred for a full neuropsychological battery [35]. On the contrary, a higher specificity may reduce the number of referrals, but many true cases could be overlooked [35]. A lower cutoff score has the potential to provide a better balance between true positives and false positive results [35]. It could also be used to identify individuals that should be assessed with a comprehensive neuropsychological battery [35].

The MoCA is accurate in detecting the presence of cognitive impairment, but it is less sensitive for determining whether it is mild or severe according to the MCCB criteria, which is the standard for assessing schizophrenia patients [35]. A baseline evaluation upon admission would enable the clinician to track cognitive changes over time, within and between admission, and identify patients that may be more vulnerable to these cognitive impairments [35].

The MoCA presents many benefits with its' use. It significantly decreases the cognitive assessment time and costs for both the patient and the provider. However, a diagnosis should not be made on MoCA results alone. A possible solution to enhance the sensitivity and specificity of the MoCA could consist of an additional short battery of tests. The MoCA can be performed in settings with limited resources, or even bedside. Minor adjustments could be made to the current administration and scoring, that could answer additional questions. For instance, the executive functioning tasks could potentially be timed to examine processing speed [35]. With further research, there may be available norms related to processing speed in different populations. The MoCA appears to have the potential to be a promising screening tool for individuals with schizophrenia, especially when a lower than the recommended threshold of 26 is used, but further research is needed in this domain [35].

### **The Impact of Illicit Drug use on MoCA Score**

Patients in an acute state of crisis are sometimes admitted to an inpatient facility with illicit drugs in their system. A drug screen via urine analysis is performed upon admission at our facility of interest. For the purposes of this study, it is important to be cautious and consider the impact that illicit drugs may have on cognitive performance, even if the patient has been hospitalized for a few days prior to testing.

Cognitive impairment in methamphetamine (MA) users with psychosis may be more severe than that in MA users without psychosis. A recent study compared the overall cognitive functioning and specific cognitive domains between MA users with and without formally assessed psychosis. Participants

included 113 MA users with psychosis and 120 MA users without psychosis. The recent use of MA, within one month prior to admission, was confirmed using quantitative analysis of hair. The MoCA total and individual domain scores were used to compare participants with and without psychosis. The association between MA psychosis and the MoCA total scores was still statistically significant after the adjustment for years in education in an ordinal logistic regression analysis. Patients who use MA with psychosis had poorer overall cognitive functioning than MA users without psychosis. Cognitive impairment was prominent in the domains of visuospatial/executive function and abstraction.

Cannabis is the most widely used illicit substance globally. A study assessed 50 young male patients who were admitted to a rehabilitation facility for cannabis use. The MoCA was used to assess the baseline cognitive functioning of these patients initially and after 1 week of abstinence from cannabis use. The patient scores were then compared with 50 graduate students that had no history of cannabis use. The MoCA was re-administered to the patient subjects after 3 months of strict abstinence from cannabis use. There was a statistically significant difference between the baseline MoCA score of cannabis users and the controls. Both the duration and the quantity of cannabis use in the patient group had a negative correlation with the MoCA score. When the MoCA test was re-administered after 3 months of abstinence, there was a statistically significant improvement in cognitive functioning in patients who used cannabis. However, the patient mean score was still less than the mean score of the student group showing only partial improvement. This study demonstrated the cognitive deficits that can be seen in cannabis users as compared to nonusers. It also showed that abstinence from cannabis in the patient sample had partially reversed the impairment, but still some deficits remained.

In sum, despite the information that a urine analysis upon admission can provide, the potential impact of illicit drug use on MoCA score is a limitation of this study.

### Study Purpose and Rationale

The overarching intention of this research project was to further understand the impact that formally assessed cognitive impairment via the Montreal Cognitive Assessment (MoCA) in psychiatric inpatients has on readmission rates. Overall, this study was designed to investigate an efficient process to assess and identify patients that may be more vulnerable to relapse and readmission.

Verbal ability, verbal memory, and executive functioning are significantly associated with problem solving in psychiatric inpatients [17]. However, cognitive functioning via formal neuropsychological assessment in inpatient settings is rarely obtained [17]. Neuropsychological testing is an underutilized resource within inpatient facilities due to cost, length of the test battery, and a limited number of neuropsychologists within these settings [36]. Implementing a neuropsychological screener can help identify deficits that may interfere with a patient's ability to adhere to a treatment plan, monitor medications, or actively participate in treatment programs; factors that are associated with readmission rates.

Increased emphasis and attention in psychiatric research have yielded more evidence to the neurobiology of cognition;

especially in Schizophrenia, Bipolar Disorder, and Post Traumatic Stress Disorder [36]. Although many psychiatric clinicians are trained in cognitive assessments, such as the "clock test," "serial sevens," "numbers forward and backward," "proverb," and "word recall," in addition to common scenarios to evaluate judgment and insight, such as "mailing a letter," they are typically not completed during a standard psychiatric evaluation due to time limitations and high volume in the unit setting [36]. Neuropsychological testing can help to pinpoint cognitive deficits that interfere with intervention and the potential for optimal patient outcomes [37-40].

This study also examined the utilization of community resources, such as outpatient treatment, medication management, housing and shelter assistance, financial aid programs, and insurance coverage. The relationship between community resource usage and readmission rates was examined. Community resource utilization data was obtained from the patient's psychosocial form archival data from a HIPAA-compliant database. Patients will be given a composite score, like the ACEs, in which each resource available will count as one as be utilized to assess return rates as a whole [40-45].

### Implications for Psychiatric Facilities, Patients, & Policy

High rates of readmission may adversely impact healthcare spending. This study may aid the composition of focused health policies to address readmission factors and improve community-based care. Incorporating a brief cognitive assessment in the evaluation of individuals with schizophrenia will improve detection of cognitive deficits and may inform treatment planning. Moreover, the results should also be corroborated with the patient's real-world functioning, for example, medication or treatment adherence, functional disability, and related outcomes. A plan that includes collaboration with other disciplines such as occupational therapy and memory-based inpatient care could either be implemented indefinitely or for a time-limited trial [46-50].

### Objectives and Hypotheses

**Objective 1:** To formally assess cognitive impairment in psychiatric inpatients via the Montreal Cognitive Assessment (MoCA) and examine the relationship between MoCA score and inpatient readmission rate.

*Hypothesis 1.1 (H1.1):* As MoCA score decreases, patient readmission rate will increase. Greater cognitive impairment as assessed by the MoCA will increase the likelihood of a patient being readmitted to an inpatient facility.

**Objective 2:** To examine the relationship between MoCA score and patient diagnosis.

*Hypothesis 1.2 (H1.2):* Patients who carry a diagnosis of a psychotic disorder will obtain lower MoCA scores.

*Hypothesis 1.3 (H1.3):* Patients who carry dual diagnoses, such as a psychiatric diagnosis in addition to a substance use disorder diagnosis, will obtain lower MoCA scores.

*Hypothesis 1.4 (H1.4):* Patients who carry a psychotic disorder diagnosis or a dual diagnosis will be younger in age, compared to the overall sample.



**Objective 3:** To examine the relationship between readmission rate and patient demographics, such as sex, age, marital status, level of education, undomiciled at the time of testing, living alone, and local family support.

*Hypothesis 1.5 (H1.5):* Readmission rate will decrease with increasing age.

*Hypothesis 1.6 (H1.6):* Readmission rate will be lower for patients that are married, and higher for those who are single, separated, or divorced.

*Hypothesis 1.7 (H1.7):* Readmission rate will be greater for patients that obtained twelve or fewer years of education.

*Hypothesis 1.8 (H1.8):* Readmission rate will be greater for patients that were undomiciled at the time of testing.

*Hypothesis 1.9 (H1.9):* Readmission rate will be higher for patients who live alone compared to those who live with others.

*Hypothesis 2.0 (H2.0):* Readmission Rate will be reduced for patients receiving local family support.

**Objective 4:** To examine the relationship between the utilization of community resources, patient readmission rate, and diagnosis, age, marital status, and education.

*Hypothesis 2.1 (H2.1):* Readmission rate will decrease with increased community resource utilization (i.e., 1 or more).

*Hypothesis 2.2 (H2.2):* Diagnosis will have a relationship with community resource usage, with individuals diagnosed with mood disorders being more likely to utilize community resources compared to those with other diagnoses.

*Hypothesis 2.3 (H2.3):* Age will be positively correlated with community resource usage, indicating that older individuals are more likely to utilize community resources.

*Hypothesis 2.4 (H2.4):* Married individuals are less likely to use community resources compared to single or divorced individuals.

*Hypothesis 2.5 (H2.5):* Higher educational attainment is positively associated with community resource usage, with individuals with more education being more likely to utilize these resources.

*Hypothesis 2.6 (H3.1):* Individuals who are homeless or living in unstable housing situations have higher community resource usage compared to those with stable housing.

## Method

This study utilized a cross-sectional design. Permission was obtained by the researcher from the Florida Institute of Technology Institutional Review Board (IRB) prior to the collection of data to utilize the cooperating inpatient psychiatric hospital's archival data. Permission was also obtained from the president of the hospital to use the archival data, which was provided to the IRB of FIT as well. The data has been entered into a HIPAA-compliant EMR database. All personal identifying information was de-identified to ensure minimal risk

of breaching confidentiality. Descriptive statistics, including assessment of means, standard deviations, and frequencies, were calculated for participant demographic variables, the primary outcomes, etc. All patient information was de-identified and random identification numbers were used for patient information on the excel sheet [51-55].

## Participants and Recruitment

Eligible participants were required to be: (1) at least 18 years of age or older, (2) have been admitted to the inpatient psychiatric facility in which the study took place and (3) were ordered a MoCA by their provider starting in January 2020 (and beyond) as it is related to increased patient protection and placement expectations utilizing assessment data. These changes took place following a review of treatment programs by the facility, which created a mandate to increase discharge placement stability through increased use of assessment data during residents' hospitalization. Of the available assessment materials, neuropsychological screening tools and the identification of potential cognitive impairment within an inpatient setting would assist with overall patient success and decision-making outcomes. The program has since utilized neuropsychological testing to inform intervention with consideration to a patient's cognitive capacity and recommend compensatory strategies to aid with medication compliance.

## Data Collection Procedures

The data that this study utilized is archival and has been entered into a HIPAA-compliant EMR database within the inpatient facility. MoCA score, total number of admissions, diagnosis, age, sex, marital status, level of education, race, insurance status, undomiciled at the time of testing, whether the patient lives alone or has local family support, medications at the time of testing, and reported community resource usage were collected from each participating patient chart. A sample of 135 patients, all of whom were given the MoCA between January 2020 and August 2023, were reviewed.

## Demographics

Demographic characteristics that were collected included diagnosis, age, sex, marital status, level of education, race, insurance status, undomiciled at the time of testing, whether the patient lives alone or has local family support, medications at the time of testing, and reported community resource usage.

## Measures

### The Montreal Cognitive Assessment (MoCA)

The Montreal Cognitive Assessment (MoCA) has been utilized with individuals in acute inpatient psychiatric settings as a screening instrument with variable results [31]. It was developed in 2005 for the purpose of detecting mild cognitive impairment (MCI) and its' results have been shown to be highly sensitive and specific in the older adult population [32]. The MoCA is a brief neuropsychological screener that is succinct enough to be administered at a patient's bedside. Total administration time typically runs about 10 to 15 minutes. The MoCA assesses short-term memory, attention, working memory, and executive functions, which are commonly affected in patients admitted to a crisis stabilization unit, or inpatient psychiatric hospitals. Scores on the MoCA range from zero to 30 points, with a score of 25 or lower indicating a cognitive impairment [31]. This cut-off score



of 25 or lower is widely recognized as a threshold for detecting cognitive impairment and potential dementia. Practice effects have been minimized as there are three versions of the MoCA, which test the same domains but differ in content [31]. The alternative versions of the MoCA present comparable reliability to the original test [33]. The MoCA has also been translated into more than 60 languages [31]. In addition, training is available on the MoCA website. Certification for the MoCA was made a mandatory requirement as of September 2019 [31]. The process of training is standardized so that testing is consistent among multiple raters, minimizing errors, and thus minimizing liability and misdiagnosis [31].

### Readmission Rate

The total admission rate per participant for a sample of 135 participants was collected. This was accessed within each patient chart, utilizing number of admissions as a metric for readmission rate. Length of stay was not a consideration for readmission, but was considered as a limitation for the number of admissions that could occur with a patient in any given year. All patient information was de-identified and random identification numbers were used for patient information on the excel sheet.

### Community Resources

Reported community resource usage was calculated by reviewing a patient's psychosocial report which is detailed as part of each patient's discharge plan. Each community resource was assigned one point. For instance, if a patient reported attending a medication management program one time per month, they would be assigned one point. If a patient reported utilizing a service that provided them with shelter, food, and a part-time job, this community resource would be assigned three points. The total community resource usage points were calculated for each participating patient and entered into an on-site excel sheet. A list of community resources is provided in Appendix A. All patient information was de-identified and random identification numbers were used for patient information on the excel sheet [56-60].

### Research Design and Data Analysis

This study utilized a cross-sectional design. The data utilized was archival and has been entered into a HIPAA-compliant EMR database within the inpatient facility. MoCA score, total number of admissions, and reported community resource usage were collected from each participating patient's chart. A sample of 135 charts of patients who were given a MoCA were reviewed.

### Statistical Analysis

Descriptive statistics, means, standard deviations, and frequencies, were calculated for participant demographics and the primary outcomes (MoCA score, readmission rate, and community resource usage score). A community resource usage score was calculated for each participant based upon what was reported in their psychosocial report. A correlation was run to examine the relationship between MoCA score, readmission rate, and total community resource usage. A linear regression was utilized to examine whether MoCA score predicts readmission rate. In addition, community resource usage score was examined as a mitigating factor to readmission rate [60-65]. Data were analyzed through the Statistical Package for the Social Sciences (SPSS), version 25.0. All analyses were considered significant at the  $p < .05$  level.

## Results

### Normality

The dataset exhibits a reassuring characteristic of normality, where the distribution of values follows a bell-shaped, symmetrical pattern. Kurtosis, a measure of the distribution's shape and propensity for extreme values, aligns with the characteristics of a standard, normal distribution. The normality in both distribution and kurtosis lends statistical robustness to the dataset, allowing for more confident and reliable analyses and inferences [66].

### Collinearity Diagnostics

In the realm of regression analysis, managing multicollinearity is crucial to ensure the reliability and integrity of results. Acceptable levels for Variance Inflation Factor (VIF) and Tolerance are often determined to strike a balance between predictor variables' intercorrelation and their influence on the model. A VIF value that is above 10 or a Tolerance value that is below .10 is problematic, as it is indicative of multicollinearity. VIF values that are below 10 or Tolerance values that are above .10, are suggestive that the variance of regression coefficients is not substantially inflated due to multicollinearity. This threshold ensures that the model remains stable and that coefficient estimates remain reasonably interpretable. In this study, the VIF and Tolerance values were examined for all four models. All VIF values were below 10 and Tolerance values were above .10, suggesting that the variance of regression coefficients was not substantially inflated due to multicollinearity [67,68].

## Participants

### Sample Demographics

A total of 135 participants were included in this study with 49.6% of participants identified as male ( $n = 67$ ) and 50.4% identified as female ( $n = 68$ ). The mean age was 69.09 years with a standard deviation of 8.25. Regarding participant ethnicity, 80.7% ( $n = 109$ ) of the participants identified as White/Caucasian, 11.9% ( $n = 16$ ) as Black, 5.2% ( $n = 7$ ) as Hispanic, and 2.2% ( $n = 3$ ) as Asian. In terms of marital status, 22% ( $n = 30$ ) reported themselves to be single, 3% ( $n = 4$ ) had a partner, 16% ( $n = 22$ ) were married, 33% ( $n = 45$ ) were divorced, 22.2% ( $n = 30$ ) were widowed, and 3% ( $n = 4$ ) were separated. The mean education level was 12.84 years with a standard deviation of 2.15. Regarding health insurance, 92.6% ( $n = 125$ ) were insured and 6.7% ( $n = 9$ ) were uninsured. In terms of housing and living situation, 23.7% ( $n = 32$ ) reported being undomiciled and 53.3 ( $n = 72$ ) lived alone. 41.5% ( $n = 56$ ) reported having local family support. Participant demographic information is presented in Table 1.

**Table 1: Descriptive Statistics of Participant Demographic Information**

Variable	N	Total Sample (N = 135) (%)
Depressive Disorders		
MDD	48	(35.6%)
MDD psychotic features	17	(12.6%)
MDD with AUD	10	(7.4%)
MDD Co-Occurring	17	(12.6%)
PDD	2	(1.5%)

Bipolar Disorders		
Bipolar 1	14	(10.4%)
Bipolar 1 Co-Occurring	2	(1.5%)
Bipolar 2	1	(1.5%)
Bipolar 2 Co-Occurring	2	(2.2%)
Psychotic Disorders		
Schizophrenia	10	(7.4%)
Schizoaffective	5	(3.7%)
Schizoaffective Co-Occurring	2	(1.5%)
Other Disorders		
Disruptive Impulse Control	1	(.7%)
Acute Stress Disorder	1	(.7%)
Co-Occurring Diagnoses		
Mental Health	56	(41.5%)
Substance Abuse	35	(25.9%)

**Descriptive Statistics**

**Diagnosis**

In terms of depressive disorders, 35.6% ( $n = 48$ ) of the sample had a diagnosis of Major Depressive Disorder, 12.6% ( $n = 17$ ) had Major Depressive Disorder with psychotic features, 7.4% ( $n = 10$ ) were given Major Depressive Disorder with Alcohol Use Disorder, and 12.6% ( $n = 17$ ) had Major Depressive Disorder with a Co-Occurring disorder (e.g., Cocaine Use Disorder or Post Traumatic Stress Disorder) [69].

10.4% ( $n = 14$ ) of the sample had received a diagnosis of Bipolar 1, 1.5% ( $n = 2$ ) had received a Bipolar 1 with a Co-Occurring Disorder, 1.5% ( $n = 2$ ) had received a diagnosis of Bipolar 2, and 2.2% ( $n = 3$ ) had received a Bipolar 2, Co-Occurring with another disorder (e.g., Cocaine Use Disorder, Cannabis Use Disorder, or Post Traumatic Stress Disorder).

Regarding psychotic disorders, 7.4% ( $n = 10$ ) had received a diagnosis of Schizophrenia, 3.7% ( $n = 5$ ) had received a schizoaffective disorder, and 1.5% ( $n = 2$ ) had received a schizoaffective disorder with a co-occurring disorder (e.g., Cocaine Use Disorder, Cannabis Use Disorder, or Post Traumatic Stress Disorder) [70].

Other diagnoses included Other Disruptive Impulse Control Disorders .7% ( $n = 1$ ), Acute Stress Disorder .7% ( $n = 1$ ), and Persistent Depressive Disorder 1.5% ( $n = 2$ ). 41.5% ( $n = 56$ ) of the sample had received a Co-Occurring diagnosis and 25.9% ( $n = 35$ ) had received a Co-Occurring substance use diagnosis. Participant diagnoses information is presented in Table 2.

**Table 2: Participant Diagnoses Information**

Variable		Total Sample (N = 135)
	M	(SD)
Age	69.09	(8.25)
	N	(%)
Gender		
Female	68	(50.4%)
Male	67	(49.6%)

Race		
White/Caucasian	109	(80.7%)
Black	16	(11.9%)
Hispanic	7	(5.2%)
Asian	3	(2.2%)
Education Level		
7 <sup>th</sup> Grade	2	(1.5%)
8 <sup>th</sup> Grade	1	(.7%)
9 <sup>th</sup> Grade	6	(4.4%)
10 <sup>th</sup> Grade	2	(1.5%)
12 <sup>th</sup> Grade	76	(56.3%)
14 Years	29	(21.5%)
16 Years	13	(9.6%)
18 Years	5	(3.7%)
21 Years	1	(.7%)
Marital Status		
Single	30	(22.2%)
Partnered	4	(3.0%)
Married	22	(16.3%)
Divorced	45	(33.3%)
Widowed	30	(22.2%)
Separated	4	(4%)
Health Insurance		
Insured	125	(92.6%)
Indigent	9	(6.7%)
Housing		
Have Housing	103	(76.3%)
Undomiciled	32	(23.7%)
Living environment		
Lived Alone	72	53.3%
Lived with Others	63	(46.7%)
Local Family Support		
Had Family Support	56	(41.5%)
Did not have Family Support	44	(58.5%)

**Montreal Cognitive Assessment (MoCA) Score**

All 135 participants completed the Montreal Cognitive Assessment. The scores ranged from a low of 2 to a high of 28. The average score for this sample was 17.20, with a standard deviation of 5.48. Participant MoCA score information is presented in Table 3.

**Table 3: MoCA Score Frequencies**

Readmission Rate	Total Sample (N = 135)	
	Frequency	Percent
2	2	1.5%
3	1	.7%
4	1	.7%
7	2	1.5%
8	2	1.5%
9	3	2.2%

10	4	3.0%
11	4	3.0%
12	7	5.2%
13	7	5.2%
14	9	6.7%
15	9	6.7%
16	5	3.7%
17	8	5.9%
18	11	8.1%
19	13	9.6%
20	9	6.7%
21	9	6.7%
22	8	5.9%
23	4	3.0%
24	3	2.2%
25	4	3.0%
26	7	5.2%
27	2	1.5%
28	1	07%

**Readmission Rate**

According to the facility’s electronic medical record system, the average readmission rate across all patients since 2014 was 1.89 with a standard deviation of 2.68. The average readmission rate for this study’s sample was 3.13 per individual with a standard deviation of 3.97. Readmission rates in this study’s sample ranged from a minimum of 1 to a high of 26 admissions. The average number of times a patient was involuntarily hospitalized for this study’s sample was 2.67 with a standard deviation of 3.27, ranging from a low of 0 to high of 26. Voluntary admissions have an average of .44 for this sample with a standard deviation of 1.49, ranging from a low of 0 to a high of 11. Information on participant readmission rate and admission type is presented in Table 4.

**Table 4: Readmission Frequencies**

Readmission Rate	Total Sample (N = 135)	
	Frequency	Percent
1	57	42.2%
2	27	20%
3	16	11.9%
4	13	9.6%
5	8	5.9%
6	2	1.5%
7	2	1.5%
8	4	3.0%
11	1	.7%
15	1	.7%
18	1	.7%
19	1	.7%
24	1	.7%
26	1	.7%

**Community Resource Usage Score**

Overall, patients in this sample reported taking advantage of an average of .96 of the Community Resources available to them, with a standard deviation of 1.89, ranging from 0 to 6. 46.7% (*n* = 63) reported utilizing no community resources at all. 29.6% (*n* = 49) reported using at least 1 community resource. 8.1% (*n* = 11) reported utilizing at least two community resources, while 13.3% (*n* = 18) used at least three. 1.5% (*n* = 2) utilized at least 4 community resources, while .7% (*n* = 1) used at least six. Patients reported utilizing outpatient medication management, outpatient therapy, forensic mental health programs and case management, nursing home services, social security, homeless shelters, soup kitchens, assisted living facilities, Alzheimer’s support groups, caregiver support groups, senior living advisors, long-term case managers, Meals on Wheels, primary care physicians, Alcoholics/Narcotics Anonymous, and Veterans’ Association Clinics [71].

**Statistical Analyses**

**MoCA Score and Readmission Rate**

A bivariate correlation was conducted to explore the relationships between the variables MoCA score, readmission rate, community resource usage, involuntary hospitalizations, and voluntary admissions. MoCA score (*M* = 17.20, *SD* = 5.49) was positively correlated with number of involuntary hospitalizations (*M* = 2.67, *SD* = 3.27), *r* = .182, *p* < .001. Total readmission rate (*M* = 3.13, *SD* = 3.97) was positively correlated with number of involuntary hospitalizations (*M* = 2.67, *SD* = 3.27), *r* = .933, *p* < .001, and voluntary admissions (*M* = 0.44, *SD* = 1.49), *r* = .615, *p* < .001. Information for MoCA Score and Readmission Rate are presented in Table 5.

**Table 5: Pearson’s Correlation Values for MoCA Score and Readmission Rate**

MoCA Score	Total Readmissions	Community Resource Usage Score	Baker Acts	Voluntary Admissions
Pearson Correlation	0.136	0.016	.182*	-0.038
Sig. (2-tailed)	0.117	0.854	0.035	0.659
Pearson Correlation		0.077	.933**	.615**
Sig. (2-tailed)		0.374	0	0
Pearson Correlation			0.041	0.111
Sig. (2-tailed)			0.636	0.201
Pearson Correlation				.290**
Sig. (2-tailed)				0.001

\*. Correlation is significant at the 0.05 level (2-tailed).

\*\*. Correlation is significant at the 0.01 level (2-tailed).

A multiple regression was conducted to examine if MoCA score predicts the number of patient readmissions to an inpatient psychiatric facility (Objective 1). MoCA score (*M* = 17.20, *SD* = 5.49) did not significantly predict patent readmission rate (*M* =



3.13, SD = 3.97),  $b = .098$ ,  $p > .001$ , which is not in supportive of Hypothesis 1.1. MoCA score did not explain a significant amount of the variance in patient readmission rate  $R^2 = .018$ ,  $F(1, 133) = 2.49$ ,  $p > .001$ . MoCA score ( $M = 17.20$ ,  $SD = 5.49$ ) also did not significantly predict the number of patient involuntary hospitalizations, or voluntary admissions ( $M = 2.67$ ,  $SD = 3.27$ ),  $b = .108$ ,  $p > .001$ . MoCA score did not explain a significant amount of the variance in the number of patient involuntary hospitalizations,  $R^2 = .033$ ,  $F(1, 133) = 42.28$ ,  $p > .001$ . MoCA score ( $M = 17.20$ ,  $SD = 5.49$ ) also does not predict the number of voluntary admissions ( $M = 0.44$ ,  $SD = 1.49$ ),  $b = -.010$ ,  $p > .001$ , which is not in support of Hypothesis 1.1. MoCA score also did not explain the amount of variance in number of patients' voluntary admission rates,  $R^2 = .001$ ,  $F(1, 133) = .437$ ,  $p > .001$ .

**MoCA Score and Diagnosis**

The average MoCA score was calculated for participants with a diagnosis of a psychotic disorder ( $n=34$ ) and was 16.26, which is lower than the overall sample mean MoCA score of 17.20. The average MoCA score was also calculated for participants with a co-occurring substance use diagnosis ( $n=33$ ) and was 17.30,

which is slightly higher than the overall sample mean MoCA score of 17.20.

A bivariate correlation was conducted to examine the relationship between MoCA score, patient diagnosis, and age (Objective 2). MoCA score ( $M = 17.20$ ,  $SD = 5.49$ ) was positively correlated with a diagnosis of Bipolar 2 Co-Occurring ( $M = .02$ ,  $SD = .15$ ),  $r = .197$ ,  $p < .001$ . MoCA score ( $M = 17.20$ ,  $SD = 5.49$ ) was negatively correlated with a diagnosis of Schizophrenia ( $M = .74$ ,  $SD = .26$ ),  $r = -.222$ ,  $p < .001$ . Age ( $M = 69.09$ ,  $SD = 8.25$ ) was positively correlated with a diagnosis of Major Depressive Disorder ( $M = .36$ ,  $SD = .48$ ),  $r = .352$ ,  $p < .001$  and a lack of a Co-Occurring Diagnosis ( $M = .59$ ,  $SD = .49$ ),  $r = -.209$ ,  $p < .001$ , which supports hypothesis 1.4. Age was also negatively correlated with readmission rate ( $M = 3.13$ ,  $SD = 3.97$ ),  $r = -.284$ ,  $p < .001$ , and a diagnosis of Bipolar 1 ( $M = .10$ ,  $SD = .31$ ),  $r = -.181$ ,  $p < .001$ . Readmission rate ( $M = 3.13$ ,  $SD = 3.97$ ) was positively correlated with Major Depressive Disorder and Alcohol Use Disorder ( $M = .074$ ,  $SD = .26$ ),  $r = .191$ ,  $p < .001$ , and Bipolar 1 disorder ( $M = .10$ ,  $SD = .31$ ),  $r = .223$ ,  $p < .001$ . Information for MoCA score and Diagnosis can be viewed in Table 6.

**Table 6: Summary of Linear Regression Analysis for MoCA Score and Total Readmissions**

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Change Statistics				
					R Square Change	F Change	df1	df2	Sig. F Change
1	.136 <sup>a</sup>	0.018	0.011	3.943	0.018	2.494	1	133	0.117

\* $p < .01$

A multiple regression was conducted to examine if a diagnosis of either a psychotic disorder or a co-occurring disorder will predict MoCA scores. A diagnosis of Major Depressive Disorder with psychotic features ( $M = .13$ ,  $SD = .33$ ),  $b = .031$ ,  $p > .001$ , Schizophrenia ( $M = .74$ ,  $SD = .26$ ),  $b = -.222$ ,  $p > .001$ , Schizoaffective ( $M = .037$ ,  $SD = .190$ ),  $b = .072$ ,  $p > .001$ , and Schizoaffective with a Co-Occurring disorder ( $M = .015$ ,  $SD = .121$ ),  $b = -.072$ ,  $p > .001$ , did not significantly predict MoCA scores ( $M = 17.20$ ,  $SD = 5.49$ ); which is not in support of Hypothesis 1.2. Additionally, a diagnosis of a psychotic disorder did not explain a significant amount of the variance in MoCA score,  $R^2 = .059$ ,  $F(1, 133) = 2.004$ ,  $p > .001$ .

The diagnoses of a co-occurring disorder did not significantly predict MoCA score which is not in support of Hypothesis 1.3. A diagnosis of Major Depressive Disorder with Alcohol Use Disorder ( $M = .074$ ,  $SD = .262$ )  $b = -.029$ ,  $p > .001$ , Major Depressive Disorder with a Co-Occurring disorder ( $M = .126$ ,  $SD = .333$ ),  $b = -.658$ ,  $p > .001$ , Bipolar 1 with a Co-Occurring disorder ( $M = .015$ ,  $SD = .121$ ),  $b = 2.871$ ,  $p > .001$ , Bipolar 2 with a Co-Occurring disorder ( $M = .022$ ,  $SD = .148$ ),  $b = 7.205$ ,  $p > .001$ , and Schizoaffective with a Co-Occurring disorder ( $M = .015$ ,  $SD = .121$ ),  $b = -3.129$ ,  $p > .001$ , did not significantly predict MoCA score ( $M = 17.20$ ,  $SD = 5.49$ ). A diagnosis of a

co-occurring disorder did not explain a significant amount of the variance in MoCA score,  $R^2 = .049$ ,  $F(1, 133) = 1.334$ ,  $p > .001$ .

**Readmission Rate and Demographic Variables**

A bivariate correlation was conducted to explore the relationships between patient demographics and readmission rates (Objective 3). Readmission rate ( $M = 3.13$ ,  $SD = 3.97$ ) was negatively correlated with African American ( $M = .118$ ,  $SD = .324$ )  $r = -.012$ ,  $p < .001$ , Hispanic, ( $M = .052$ ,  $SD = .223$ )  $r = -.016$ ,  $p < .001$ , and Asian ( $M = .222$ ,  $SD = .148$ )  $r = -.081$ ,  $p < .001$ . Readmission rate ( $M = 3.13$ ,  $SD = 3.97$ ) was also negatively correlated with Age ( $M = 69.09$ ,  $SD = 8.25$ )  $r = -.284$ ,  $p < .001$ , which is in support of Hypothesis 1.5. Being Female ( $M = .504$ ,  $SD = .502$ )  $r = -.163$ ,  $p < .001$ , was negatively correlated with readmission rate, in addition to having family support ( $M = .415$ ,  $SD = .495$ )  $r = -.179$ ,  $p < .001$ , which is in support of Hypothesis 2.0. Readmission rate was positively correlated with having a partner, but not being married or single, ( $M = .163$ ,  $SD = .371$ )  $r = .193$ ,  $p < .001$ , which is not in support of Hypothesis 1.6. Hospitalization rates were also higher for patients that were living alone ( $M = .533$ ,  $SD = .501$ )  $r = .184$ ,  $p < .001$ , at the time of admission which is in support of Hypothesis 1.9. Information on readmission rate and patient demographics can be viewed in Table 7.

**Table 7: Pearson’s Correlation Values for MoCA Score and Patient Diagnosis**

	MoCA Score	Total Readmissions	Age	YES Co-Occurring Diagnoses	Major Depressive Disorder (MDD)	MDD with Psychotic Features	MDD with Alcohol Use Disorder (AUD)	MDD with Co-Occurring Disorder	Bipolar 1	Bipolar 1 with Co-Occurring Disorder	Bipolar 2	Bipolar 2 with Co-Occurring Disorder	Schizophrenia	Schizoaffective	Schizoaffective with Co-Occurring Disorder	Other Disruptive Impulse Control Disorder	Acute Stress Disorder	Persistent Depressive Disorder	Yes Substance Use Diagnoses
1	Pearson Correlation	0.136	-0.095	-0.020	-0.036	0.031	-0.005	-0.051	0.121	0.063	0.130	.197*	-.222**	0.072	-0.072	-0.082	-0.114	0.007	0.025
	Sig. (2-tailed)	0.117	0.273	0.820	0.681	0.721	0.953	0.560	0.163	0.469	0.133	0.022	0.010	0.408	0.408	0.343	0.189	0.938	0.776
2	Pearson Correlation		-.284**	-.179*	-0.157	-0.052	.191*	0.011	.223**	-0.035	-0.050	-0.018	-0.009	-0.046	-0.035	-0.046	-0.046	0.043	0.088
	Sig. (2-tailed)		0.001	0.038	0.069	0.552	0.026	0.904	0.009	0.687	0.561	0.840	0.917	0.597	0.687	0.592	0.592	0.623	0.310
3	Pearson Correlation			.209*	.352**	-0.107	0.062	-0.053	-.181*	-0.024	-0.054	-0.142	-0.062	-0.098	-0.068	0.052	0.010	-0.039	-0.085
	Sig. (2-tailed)			0.015	0.000	0.215	0.473	0.541	0.036	0.785	0.537	0.100	0.478	0.260	0.430	0.552	0.912	0.656	0.329
4	Pearson Correlation				.191*	-0.048	-.181*	-0.093	0.009	0.021	0.021	-0.025	-0.123	0.153	0.021	-0.073	0.103	-0.103	-.189*
	Sig. (2-tailed)				0.026	0.583	0.036	0.283	0.913	0.807	0.807	0.774	0.154	0.076	0.807	0.402	0.236	0.233	0.028
5	Pearson Correlation					-.282**	-.210*	-.282**	-.253**	-0.091	-0.091	-0.112	-.210*	-0.146	-0.091	-0.064	-0.064	-0.091	-.439**
	Sig. (2-tailed)					0.001	0.014	0.001	0.003	0.293	0.293	0.196	0.014	0.092	0.293	0.460	0.460	0.293	0.000
6	Pearson Correlation						-0.107	-0.144	-0.129	-0.047	-0.047	-0.057	-0.107	-0.074	-0.047	-0.033	-0.033	-0.047	-.225**
	Sig. (2-tailed)						0.215	0.096	0.136	0.592	0.592	0.510	0.215	0.391	0.592	0.706	0.706	0.592	0.009
7	Pearson Correlation							-0.107	-0.096	-0.035	-0.035	-0.043	-0.080	-0.055	-0.035	-0.024	-0.024	-0.035	.478**
	Sig. (2-tailed)							0.215	0.267	0.690	0.690	0.623	0.356	0.523	0.690	0.778	0.778	0.690	0.000
8	Pearson Correlation								-0.129	-0.047	-0.047	-0.057	-0.107	-0.074	-0.047	-0.033	-0.033	-0.047	.642**
	Sig. (2-tailed)								0.136	0.592	0.592	0.510	0.215	0.391	0.592	0.706	0.706	0.592	0.000
9	Pearson Correlation									-0.042	-0.042	-0.051	-0.096	-0.067	-0.042	-0.029	-0.029	-0.042	-.201*
	Sig. (2-tailed)									0.631	0.631	0.555	0.267	0.442	0.631	0.735	0.735	0.631	0.019
10	Pearson Correlation										-0.015	-0.018	-0.035	-0.024	-0.015	-0.011	-0.011	-0.015	.207*
	Sig. (2-tailed)										0.863	0.831	0.690	0.782	0.863	0.903	0.903	0.863	0.016
11	Pearson Correlation											-0.018	-0.035	-0.024	-0.015	-0.011	-0.011	-0.015	-0.073
	Sig. (2-tailed)											0.831	0.690	0.782	0.863	0.903	0.903	0.863	0.403
12	Pearson Correlation												-0.043	-0.030	-0.018	-0.013	-0.013	-0.018	.255**
	Sig. (2-tailed)												0.623	0.734	0.831	0.881	0.881	0.831	0.003
13	Pearson Correlation													-0.055	-0.035	-0.024	-0.024	-0.035	-0.167
	Sig. (2-tailed)													0.523	0.690	0.778	0.778	0.690	0.052
14	Pearson Correlation														-0.024	-0.017	-0.017	-0.024	-0.027
	Sig. (2-tailed)														0.782	0.845	0.845	0.782	0.760
15	Pearson Correlation															-0.011	-0.011	-0.015	.207*
	Sig. (2-tailed)																		0.000





in turn, improve cognitive functioning. Patients who lack a co-occurring diagnosis may have a clearer clinical presentation, making it easier to identify and address primary mental health concerns [1,13]. This may lead to better treatment outcomes and cognitive functioning. Patients without co-occurring diagnoses may experience fewer overlapping symptoms. They may receive targeted treatment and psychoeducation, addressing the specific symptoms and cognitive impairments, potentially resulting in higher MoCA scores.

Lower MoCA scores were associated with diagnoses of Bipolar 1 and Schizophrenia, which may have severe courses that can be associated with cognitive deficits. Cognitive impairments can impact attention, memory, executive functioning, and processing speed, which may result in lower MoCA scores [1,13]. Bipolar 1 and Schizophrenia have been linked to neurobiological abnormalities such as alterations in ventricle size and neurotransmitter functioning [1,13]. These biological factors can contribute to cognitive dysfunction. In addition, the severity of symptoms and the frequency of mood or psychotic episodes can impact cognitive functioning. During manic or psychotic episodes, patients with Bipolar 1 and Schizophrenia may experience cognitive disorganization and impairment. The negative symptoms of schizophrenia, in particular, can be associated with cognitive deficits as they impact motivation and engagement in cognitive tasks [1,13]. During periods of euthymia, patients may perform better on cognitive assessments compared to when they are manic or severely depressed. A patient's diagnosis and present symptoms may not only affect their ability to manage their own symptoms, but they may also impact the results of their cognitive testing. The MoCA is often thought of as a test that can be conducted anytime, even at a patient's bedside. However, appropriate timing in cognitive testing is important and must be considered in the interpretation of the results.

Both Bipolar 1 and Schizophrenia are often chronic conditions with a lifelong course. Prolonged exposure to the symptoms and stress associated with these disorders may potentially have a cumulative negative impact on cognitive functioning. Some patients with these diagnoses may not have received appropriate treatment or may have limited access to mental healthcare, as demonstrated earlier in this discussion. Untreated or undertreated serious mental illness may have a relationship with severe cognitive impairment. Also, other medical and psychiatric comorbidities can further contribute to cognitive deficits. Lastly, sometimes access to community resources can be compounding; for example, if a patient is unable to obtain their medication, they may not find access to supplied housing and/or feel comfortable attending follow-up appointments.

Readmission rate demonstrated a positive relationship with Major Depressive Disorder Alcohol Use Disorder, and Bipolar 1 disorder. Severe depression, manic episodes, or alcohol withdrawal can lead to acute crises that necessitate inpatient treatment. The positive relationship between readmission rate, Major Depressive Disorder Alcohol Use Disorder, and Bipolar 1 disorder reflects the complex and multifaceted nature of these conditions. Acute episodes, crisis management needs, safety concerns, and challenges associated with co-occurring disorders all contribute to the likelihood of readmissions.

Readmission rate was negatively correlated with being African American, Hispanic, and Asian. This is a representation of the lack of diversity in this primarily Caucasian sample. Being Female was negatively correlated with readmission rate which may demonstrate a gender difference in seeking and engaging in mental health treatment, social support networks, and access to care. Gender is just one of the many factors that influence mental health outcomes and individual experiences can vary widely.

Having local family support was also negatively correlated with readmission rate. Family support can provide emotional, practical, and social support to individuals with mental health conditions. Social support can help individuals cope with stressors, manage their symptoms, adhere to their medication regimen, and prevent crises that may lead to readmissions. Family members who are actively involved in a patient's mental health care can often recognize the early signs of relapse or worsening symptoms. They can help intervene early, potentially avoiding the need for inpatient readmission. Families can act as advocates for their loved ones within the healthcare system as they can communicate with treatment providers as historians or share information about patient progress. Families can also work with mental health providers to develop safety plans and crisis intervention strategies, potentially mitigating the occurrence of inpatient readmission. Family support can also reduce feelings of isolation and disconnectedness, which are risk factors for suicide.

Readmission rate was positively correlated with having a partner, but not being married or single. The nature and dynamics of a relationship can significantly impact an individual's mental health. Having a partner, even if not legally married, can introduce unique stressors, conflict, or support dynamics that may affect mental health and contribute to readmission rate.

Hospitalization rates were also higher for patients that were living alone. Individuals who live alone but may have limited access to immediate support during times of crisis, or when experiencing severe mental health symptoms. The absence of someone who can provide intervention, assistance, or seek help may lead to a higher likelihood of hospitalization. People who live alone may also delay seeking help for mental health concerns due to a lack of immediate support or encouragement to seek treatment. Delayed intervention can result in symptoms worsening and need for hospitalization. Lastly, living alone comes with the potential for associated financial pressure, loneliness, and a lack of social support.

Community resource usage was negatively correlated with having a partner, but not being married or single. The nature of the relationship and the dynamics within it can influence an individual's willingness and ability to seek out community resources. These results suggest an intriguing dynamic within the context of community support. It may indicate that individuals in committed relationships (having a partner) already have access to a built-in support system, thereby reducing their reliance on external community resources. On the other hand, those who are single or married may find it necessary to seek support from the broader community due to the absence of or different nature of support within their relationship status. This finding underscores the complex interplay between personal

relationships and community resources in meeting individuals' needs. It highlights the importance of tailoring support systems to diverse relationship statuses for more effective community interventions.

**Table 8: Summary of Linear Regression Analysis for MoCA Score and Psychotic Disorders**

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Change Statistics				
					R Square Change	F Change	df1	df2	Sig. F Change
1	.136 <sup>a</sup>	0.018	0.011	3.943	0.018	2.494	1	133	0.117

\*p<.01

**Table 9: Summary of Linear Regression Analysis for MoCA Score and Co-Occurring Diagnoses**

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Change Statistics				
					R Square Change	F Change	df1	df2	Sig. F Change
1	.136 <sup>a</sup>	0.018	0.011	3.943	0.018	2.494	1	133	0.117

\*p<.01

**Table 10: Pearson's Correlation Values for Readmission Rate and Demographic Variables**

	Total Readmissions	Age	Education	Male	Female	Insured	Uninsured	Undomiciled	Has Housing	Lives Alone	Lives with Others	Single	Partner	Married	Divorced	Widowed	Separated	Caucasian	African American	Hispanic	Asian	Has Family Support	Not Have Family Support	
1 Pearson Correlation																								
Sig. (2-tailed)																								
2 Pearson Correlation																								
Sig. (2-tailed)																								
3 Pearson Correlation																								
Sig. (2-tailed)																								
4 Pearson Correlation																								
Sig. (2-tailed)																								
5 Pearson Correlation																								
Sig. (2-tailed)																								
6 Pearson Correlation																								
Sig. (2-tailed)																								
7 Pearson Correlation																								
Sig. (2-tailed)																								
8 Pearson Correlation																								
Sig. (2-tailed)																								
9 Pearson Correlation																								
Sig. (2-tailed)																								
10 Pearson Correlation																								
Sig. (2-tailed)																								
11 Pearson Correlation																								
Sig. (2-tailed)																								





**Table 12: Pearson's Correlation Values for Community Resource Usage and Patient Demographics**

		Age	Educational	Male	Female	Insured	Uninsured	Undomiciled	Has Housing	Lives Alone	Lives with Others	Single	Partner	Married	Divorced	Widowed	Separated	Caucasian	African American	Hispanic	Asian	Has Family Support	Do not have Family Support	Community Resource Usage Score
1	Pears on Correl Sig. (2-tailed)		-0.104	-0.049	0.049	.273	-.273	-.290	.290	-0.077	0.077	-.214	0.104	0.022	-0.092	.261	-0.013	0.133	-0.082	-0.108	-0.014	.209	-.209	.183
2	Pears on Correl Sig. (2-tailed)			0.576	0.576	0.001	0.001	0.001	0.001	0.377	0.377	0.013	0.228	0.799	0.290	0.002	0.885	0.123	0.344	0.211	0.873	0.015	0.015	0.033
3	Pears on Correl Sig. (2-tailed)			0.145	-0.145	-0.061	0.061	-0.128	0.128	-0.057	0.057	0.032	0.136	0.090	-0.130	0.032	-0.130	.209	-.240	-0.107	0.129	0.078	-0.078	0.077
4	Pears on Correl Sig. (2-tailed)			0.094	0.094	0.482	0.482	0.138	0.138	0.509	0.509	0.709	0.116	0.300	0.134	0.709	0.134	0.015	0.005	0.216	0.136	0.368	0.368	0.377
5	Pears on Correl Sig. (2-tailed)			-1.000	-0.149	0.149	0.109	0.109	-0.109	0.127	-0.127	.182	0.001	-0.117	0.052	-0.139	0.001	-0.004	0.003	0.035	-0.049	-.174	.174	-0.057
6	Pears on Correl Sig. (2-tailed)			0.000	0.086	0.086	0.210	0.210	0.143	0.143	0.034	0.988	0.176	0.546	0.109	0.988	0.967	0.967	0.975	0.686	0.571	0.043	0.043	0.514
7	Pears on Correl Sig. (2-tailed)			0.149	-0.149	-0.109	0.109	-0.127	0.127	-.182	-0.001	0.117	-0.052	0.139	-0.001	0.004	-0.003	-0.035	0.049	.174	-.174	.057		
8	Pears on Correl Sig. (2-tailed)			0.086	0.086	0.210	0.210	0.143	0.143	0.034	0.988	0.176	0.546	0.109	0.988	0.967	0.967	0.975	0.686	0.571	0.043	0.043	0.514	
9	Pears on Correl Sig. (2-tailed)			-1.000	-.339	-.339	.339	-.189	.189	-.189	.189	-0.142	0.047	0.116	0.001	0.073	-0.128	0.019	0.007	-0.071	0.041	0.163	-0.163	0.069
10	Pears on Correl Sig. (2-tailed)			0.000	0.000	0.000	0.029	0.029	0.102	0.102	0.589	0.183	0.987	0.405	0.140	0.826	0.937	0.415	0.641	0.059	0.059	0.430		
11	Pears on Correl Sig. (2-tailed)			.339	-.339	-.189	.189	-.189	.189	-.189	.189	0.142	-0.047	-0.116	-0.001	-0.073	0.128	-0.019	-0.007	0.071	-0.041	-0.163	0.163	-0.069
12	Pears on Correl Sig. (2-tailed)			0.000	0.000	0.029	0.029	0.102	0.102	0.589	0.183	0.987	0.405	0.140	0.826	0.937	0.415	0.641	0.059	0.059	0.430			
13	Pears on Correl Sig. (2-tailed)			-1.000	.452	-.452	.163	-0.097	-.199	0.086	-0.005	-0.097	-0.037	0.011	0.105	-0.084	-.399	.399	-.130					
14	Pears on Correl Sig. (2-tailed)			0.000	0.000	0.000	0.059	0.261	0.021	0.320	0.957	0.261	0.670	0.898	0.224	0.333	0.000	0.000	0.133					
15	Pears on Correl Sig. (2-tailed)			.452	-.452	-.163	0.097	.199	-0.086	0.005	0.097	0.037	-0.011	-0.105	0.084	-.399	.399	-.130						
16	Pears on Correl Sig. (2-tailed)			0.000	0.000	0.059	0.261	0.021	0.320	0.957	0.261	0.670	0.898	0.224	0.333	0.000	0.000	0.133						
17	Pears on Correl Sig. (2-tailed)			-1.000	0.071	-0.012	-.351	.220	0.000	-0.012	0.146	-0.070	-0.049	-0.161	-.478	.478	-.017							
18	Pears on Correl Sig. (2-tailed)			0.000	0.410	0.893	0.000	0.010	1.000	0.893	0.092	0.417	0.572	0.062	0.000	0.000	0.847							
19	Pears on Correl Sig. (2-tailed)			-0.071	0.012	.351	-.220	0.000	0.012	-0.146	0.070	0.049	0.161	-.478	.478	-.017								
20	Pears on Correl Sig. (2-tailed)			0.410	0.893	0.000	0.010	1.000	0.893	0.092	0.417	0.572	0.062	0.000	0.000	0.847								
21	Pears on Correl Sig. (2-tailed)			-0.093	-.236	-.378	-.286	-0.093	-0.055	0.135	-0.045	-0.081	-.305	.305	-.062									
22	Pears on Correl Sig. (2-tailed)			0.281	0.006	0.000	0.001	0.281	0.525	0.119	0.607	0.353	0.000	0.000	0.475									
23	Pears on Correl Sig. (2-tailed)			-0.077	-0.124	-0.093	-0.031	0.085	-0.064	-0.041	-0.026	0.119	-0.119	0.153										
24	Pears on Correl Sig. (2-tailed)			0.374	0.153	0.281	0.725	0.325	0.460	0.638	0.762	0.170	0.170	0.076										
25	Pears on Correl Sig. (2-tailed)			-0.312	-.236	-0.077	-0.039	-0.100	0.168	0.070	.483	-.483	-.190											
26	Pears on Correl Sig. (2-tailed)			0.000	0.006	0.374	0.655	0.250	0.051	0.423	0.000	0.000	0.028											
27	Pears on Correl Sig. (2-tailed)			.378	-.124	-0.013	0.081	-0.094	0.000	-0.117	0.117	0.142												
28	Pears on Correl Sig. (2-tailed)			0.000	0.153	0.878	0.350	0.276	1.000	0.177	0.177	0.101												
29	Pears on Correl Sig. (2-tailed)			-0.093	0.080	-0.086	-0.045	0.040	-0.052	0.052	-0.059													
30	Pears on Correl Sig. (2-tailed)			0.281	0.354	0.323	0.607	0.643	0.547	0.547	0.500													
31	Pears on Correl Sig. (2-tailed)			-0.025	-0.064	0.156	-0.026	0.030	-0.030	-0.142														
32	Pears on Correl Sig. (2-tailed)			0.770	0.460	0.071	0.762	0.728	0.728	0.100														
33	Pears on Correl Sig. (2-tailed)				-.751	-.479	-.309	.106	-0.106	0.064														
34	Pears on Correl Sig. (2-tailed)			0.000	0.000	0.000	0.220	0.220	0.460															
35	Pears on Correl Sig. (2-tailed)			-0.086	-0.055	-.216	.216	.011																
36	Pears on Correl Sig. (2-tailed)			0.323	0.524	0.012	0.012	0.895																
37	Pears on Correl Sig. (2-tailed)			-0.035	0.007	-0.007	-0.134																	
38	Pears on Correl Sig. (2-tailed)			0.685	0.940	0.940	0.121																	
39	Pears on Correl Sig. (2-tailed)			.179	-.179	.005																		
40	Pears on Correl Sig. (2-tailed)			0.038	0.038	0.957																		

20	Pears on Correl Sig. (2-tailed)																		.179	-.179	0.005
																			0.038	0.038	0.957
21	Pears on Correl Sig. (2-tailed)																			-1.000*	0.001
																				0.000	0.991
22	Pears on Correl Sig. (2-tailed)																				-0.001
																					0.991

\*. Correlation is significant at the 0.05 level (2-tailed)  
 \*\*. Correlation is significant at the 0.01 level (2-tailed)

**Table 13: Summary of Linear Regression Analysis for Patient Demographics and Community Resource Usage**

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Change Statistics				
					R Square Change	F Change	df1	df2	Sig. F Change
1	.452 <sup>a</sup>	0.205	0.103	3.765	0.205	2.023	15	118	0.019

\*p<.01

**Limitations**

One of the major limitations of this retrospective study was the reliance on an archival dataset, which hindered the range of available information specific to each participant included in the analysis. Thus, the total sample size was limited to patient data in which the MoCA had already been completed. In addition, community resource usage scores could be computed only from what was reported by each patient and recorded in the medical chart. This may mean that the community resource scores may not be exact as some patients may not have reported their usage accurately, or usage may not have been recorded appropriately.

Regarding participant ethnicity, 80.7% (n = 109) of the participants identified as White/Caucasian. A lack of patient diversity is a limitation of this study. The need for a more diverse sample to achieve representativeness is paramount in research and data analysis. A diverse sample ensures that the findings and conclusions drawn from the study are more broadly applicable to the population of interest. The results of this study lack generalizability.

Although power analyses deemed the number of participants for this study as adequate; the sample size remained small. Working with a small sample size comes with inherent limitations that can impact the reliability and generalizability of findings. The most significant limitation is reduced statistical power, which increases the risk of Type II errors, meaning that true effects can be overlooked. Additionally, this sample does not adequately represent the diversity and variability present in the larger population, limiting the external validity of the study. This study lacks robust conclusions. Subgroup differences were difficult to assess and the exploration of complex relationships between variables was hindered by the small sample size.

The orders for neuropsychological testing were mostly completed on one of the hospital’s buildings due to the nature of the cognitive concerns. This building services primarily insured, older individuals. Regarding health insurance, 92.6% (n = 125) were insured and 6.7% (n = 9) were uninsured. The facility also has a crisis intervention unit which services primarily uninsured, younger, and sometimes undomiciled patients. The addition of

uninsured, younger, and undomiciled individuals would diversify the study’s demographic composition, better reflecting a broader cross-section of society. This diversity in the sample population could have led to a more comprehensive understanding of the research topic and enhanced the study’s external validity.

The impact of psychotropic medication and illicit drug use on MoCA score is unknown and a limitation of this study. 100% of the sample was taking a psychotropic medication regimen at the time of testing. No participant in the sample were documented as using illicit drugs at the time of testing, however, there may be cognitive impacts from a drug use history; including substance use as reason for admission and occurring up until their hospitalization. While it is well recognized that medications and illicit drugs have the potential to impact cognitive functioning, the precise nature and extent of these effects on MoCA score in this study is unknown, can vary widely between individuals, and between different medications. Similarly, illicit drug use encompasses a broad spectrum of substances, each with distinct cognitive implications. The degree of cognitive impairment can depend on factors such as frequency, duration, and dose of the substance, in addition to variability within the individual patient. Therefore, while the MoCA has the potential to provide valuable insights into cognitive functioning, a comprehensive understanding of how medication and prior illicit drug use influence cognitive abilities necessitates a more nuanced examination which is more in-depth than what can be collected from an archival dataset.

**Areas for Future Research**

Future research should be longitudinal and investigate cognitive changes over time in psychiatric inpatients who are regularly readmitted. This can be accomplished with several alternative versions of the MoCA or the Repeatable Battery for the Assessment of Neuropsychological Status (R-BANS). Longitudinal research can provide insights into the trajectory of cognitive functioning during hospitalization and after discharge, helping in the development of targeted interventions. It may also help to examine factors influencing cognitive decline and recovery in psychiatric patients. This research could identify predictors of cognitive improvement, potentially guiding treatment

strategies. Upcoming longitudinal research should also study cognitive changes in patients during the transition from inpatient to community-based care. This transition may impact cognitive functioning and the risk of readmission.

Future studies should investigate the different domains of the MoCA by diagnosis to see if cognitive deficits are more generalized or follow domain-specific patterns and have a neurocognitive profile. In addition, the MoCA should be normed for different groups of patients, such as patients with schizophrenia and bipolar disorder. As discussed earlier in this study, the MoCA was utilized in diagnosing cognitive impairment in psychiatric inpatients [34]. A MoCA threshold score of 23 was the least accurate in terms of differentiating neurocognitive disorders from psychotic disorders [34]. Future research that seeks to norm the MoCA for different patient groups should also investigate cut scores that represent good sensitivity and specificity for each grouping. This research may help to develop and test cognitive rehabilitation interventions tailored to the needs of psychiatric inpatients. Additionally, the effectiveness of current interventions can be assessed and advanced, potentially improving cognitive function and overall outcomes.

Upcoming research should also investigate the impact of psychiatric medications on cognitive functioning. It should explore and seek to define how different drug classes affect cognitive performance and whether medication adjustments can mitigate cognitive side effects.

Studies could also investigate how psychosocial factors such as social support, stress, being undomiciled, and trauma history impact cognitive function in psychiatric inpatients. This research could shed light on the interplay between mental health and cognitive performance. Cultural and ethnic variations in cognitive assessment scores among psychiatric inpatients should be examined. This could highlight potential biases in assessment tools and lead to more culturally sensitive approaches.

Readmission rate or patient recidivism requires a universal definition for research purposes. It may be helpful to examine the relationship between readmission rate and the length of patient stay in an inpatient facility. As demonstrated by this study, readmission rate to psychiatric facilities is a multifaceted issue influenced by a multitude of factors, and not always attributable to the shortcomings on the part of the facility. Patients grappling with mental health challenges often face complex and dynamic circumstances which occur outside of the facility and are quite literally outside of the control of healthcare providers. As demonstrated by this study, these factors include the availability and accessibility of community mental health resources, the stability of a patient's social support system, the effectiveness of outpatient treatment plans, and individual patient factors such as diagnosis chronicity, co-occurring disorders, social support, and life-stressors. Furthermore, readmission can sometimes be a necessary and planned part of a patient's recovery journey, as evidenced by the argument which sought to explain the positive relationship between MoCA score and readmission rate. Inpatient facilities play a crucial role in our healthcare system. However, reducing readmission rates requires a comprehensive approach that addresses a broader ecosystem of factors, which impacts the well-being of individuals living with mental health conditions.

Each of these future research directions has the potential to enhance our understanding of cognitive functioning in inpatient psychiatric populations, inform clinical practice, and contribute to improving the quality of care and outcomes for individuals with mental health disorders.

### Clinical Implications

There are many ways that this research informs clinical practice. While a higher readmission rate to a psychiatric facility being associated with a higher MoCA score may seem paradoxical, multiple factors, including the severity of mental illness, reasons for readmission, and patient characteristics contribute to this study's results. Tailoring treatment and rehabilitation strategies to individual cognitive profiles may enhance clinical outcomes. Understanding the reasons for readmission is essential. Identifying these reasons can inform targeted interventions that may reduce the risk of readmission. Individual patient characteristics, such as age, co-occurring conditions, and psychosocial factors, play a significant role in cognitive function. Clinicians should conduct comprehensive assessments that consider these factors to better understand cognitive performance and its relationship to readmission risk. A nuanced approach to cognitive assessment is warranted.

The association between higher readmission rates and higher MoCA scores in psychiatric patients highlights the need for individualized and multifaceted assessments and interventions. Clinicians must consider the multifactorial nature of cognitive function in this population and strive to address the complex interplay of mental illness severity, readmission reasons, and patient characteristics to optimize patient care and outcomes.

Assessing cognitive deficits within psychiatric settings may play a pivotal role in understanding and addressing the complex issue of readmission rates. While demographic factors and limited access to community resources are undoubtedly crucial considerations, evaluating cognitive functioning adds an extra layer of insight that may have been traditionally overlooked. Cognitive deficits, often linked to conditions like schizophrenia, bipolar disorder, or major depressive disorder, can significantly impact an individual's ability to manage their mental health effectively. Patients with impaired cognitive functioning may struggle with medication adherence, coping skills, and daily life tasks, ultimately increasing the risk of relapse and readmission. By identifying and addressing cognitive deficits alongside demographic and resource-related challenges, psychiatric healthcare providers can tailor interventions and support services more precisely, improving the overall quality of care and reducing the likelihood of recurrent hospitalizations.

### References

1. Bowie CR, Harvey PD. Cognitive deficits and functional outcome in schizophrenia. *Neuropsychiatric Disease and Treatment*. 2006 2: 531-536.
2. Davis DH, Creavin ST, Yip JL, Noel-Storr AH, Brayne C, et al. Montreal Cognitive Assessment for the diagnosis of Alzheimer's disease and other dementias. *Cochrane Database of Systematic Reviews*. 2015. 10.
3. American Psychiatric Association. Practice Guideline for the Assessment and Treatment of Patients with Suicidal Behaviors. 2003. 30-33.



4. National Mental Health Services Survey. US Department of Health and Human Services, National Institutes of Health, National Institute of Mental Health. 2020. [https://www.samhsa.gov/data/sites/default/files/reports/rpt35336/2020\\_NMHSS\\_final.pdf](https://www.samhsa.gov/data/sites/default/files/reports/rpt35336/2020_NMHSS_final.pdf)
5. Shah B, Leontieva L, Megna JL. Shifting Trends in Admission Patterns of an Acute Inpatient Psychiatric Unit in the State of New York. *Cureus*. 2020. 12.
6. Wadley VG, Okonkwo O, Crowe M, Ross-Meadows LA. Mild cognitive impairment and everyday function: Evidence of reduced speed in performing instrumental activities of daily living. *American Journal of Geriatric Psychiatry*. 2008. 16: 416-424.
7. Borson S. Cognition, aging, and disabilities: conceptual issues. *Physical Medicine and Rehabilitation Clinics of North America*. 2010. 21: 375-382.
8. Kim A. The Montreal Cognitive Test Intervention, Professional Case Management. 2020. 25: 85-91.
9. Lyons JS, O'Mahoney MT, Miller SI, Neme J, Kabat J, et al. Predicting readmission to the psychiatric hospital in a managed care environment: implications for quality indicators. *The American journal of psychiatry*. 154: 337-340.
10. Fisekovic S, Memic A, Pasalic A. Correlation between moca and mmse for the assessment of cognition in schizophrenia. *Acta informatica medica : AIM : journal of the Society for Medical Informatics of Bosnia & Herzegovina : casopis Drustva zamedicinsku informatiku BiH*. 20: 186-189.
11. Bilder RM, Goldman RS, Volavka J, Czobor P, Hoptman M, et al. Neurocognitive effects of clozapine, olanzapine, risperidone, and haloperidol in patients with chronic schizophrenia or schizoaffective disorder. *The American journal of psychiatry*. 2002. 159: 1018-1028.
12. Cornblatt BA, Erlenmeyer-Kimling L. Global attentional deviance as a marker of risk for schizophrenia: specificity and predictive validity. *Journal of abnormal psychology*. 1985. 94: 470-486.
13. Caspi A, Reichenberg A, Weiser M, Rabinowitz J, Kaplan Z, et al. Cognitive performance in schizophrenia patients assessed before and following the first psychotic episode. *Schizophrenia research*. 2003. 65: 87-94.
14. Kerns JG, Berenbaum H, Barch DM, Banich MT, Stolar N. Word production in schizophrenia and its relationship to positive symptoms. *Psychiatry research*. 1999. 87: 29-37.
15. Aloia MS, Gourovitch ML, Weinberger DR, Goldberg TE. An investigation of semantic space in patients with schizophrenia. *Journal of the International Neuropsychological Society: JINS*. 1996. 2: 267-273.
16. Goldberg TE, Aloia MS, Gourovitch ML, Missar D, Pickar D, et al. Cognitive substrates of thought disorder, I: the semantic system. *The American journal of psychiatry*. 1998. 155: 1671-1676.
17. Addington J, Addington D. Neurocognitive and social functioning in schizophrenia: a 2.5 year follow-up study. *Schizophrenia research*. 2000. 44: 47-56.
18. McGurk SR, Coleman T, Harvey PD, Reichenberg A, White L, et al. Working memory performance in poor outcome schizophrenia: relationship to age and executive functioning. *Journal of clinical and experimental neuropsychology*. 2004. 26: 153-160.
19. Mausbach BT, Bowie CR, Harvey PD, Twamley EW, Goldman SR, et al. Usefulness of the UCSD performance-based skills assessment (UPSA) for predicting residential independence in patients with chronic schizophrenia. *Journal of psychiatric research*. 2008. 42: 320-327.
20. Rosca P, Bauer A, Grinshpoon A, Khawaled R, Mester R, et al. Rehospitalizations among psychiatric patients whose first admission was involuntary: A 10-year follow-up. *Isr. J. Psychiatry Relat. Sci*. 2006. 43: 57.
21. Durbin J, Lin E, Layne C, Teed M. Is readmission a valid indicator of the quality of inpatient psychiatric care? *J. Behav. Health Serv. Res*. 2007. 34: 137-150.
22. Bozikas VP, Kosmidis MH, Karavatos A. Disproportionate impairment in semantic verbal fluency in schizophrenia: Differential deficit in clustering. *Schizophrenia Research*. 74: 51-59.
23. Collins AG, Brown JK, Gold JM, Waltz JA, Frank MJ. Working memory contributions to reinforcement learning impairments in schizophrenia. *Journal of Neuroscience*. 34: 13747-13756.
24. Lepage M, Bodnar M, Bowie CR. Neurocognition: clinical and functional outcomes in schizophrenia. *The Canadian Journal of Psychiatry*. 2014. 59: 5-12.
25. Wu C, Dagg P, Molgat C. A pilot study to measure cognitive impairment in patients with severe schizophrenia with the Montreal Cognitive Assessment (MoCA). *Schizophrenia research*. 2014. 158: 151-155.
26. Désaméricq G, Schurhoff F, Meary A, Szöke A, Macquin-Mavier I, et al. Long-term neurocognitive effects of antipsychotics in schizophrenia: a network meta-analysis. *European journal of clinical pharmacology*. 2014. 70: 127-134.
27. Ravaglia G, Forti P, Maioli F, Servadei L, Martelli M, et al. Screening for mild cognitive impairment in elderly ambulatory patients with cognitive complaints. *Aging clinical and experimental research*. 2005. 17: 374-379.
28. Anderson TM, Sachdev PS, Brodaty H, Trollor JN, Andrews G. Effects of sociodemographic and health variables on Mini-Mental State Exam scores in older Australians. *The American Journal of Geriatric Psychiatry*. 2007 15: 467-476.
29. Mackin RS, Ayalon L, Feliciano L, Areán PA. The sensitivity and specificity of cognitive screening instruments to detect cognitive impairment in older adults with severe psychiatric illness. *Journal of geriatric psychiatry and neurology*. 2010. 23: 94-99.
30. Cullum CM, Saine K, Chan LD, Martin-Cook K, Gray KF, et al. Performance-Based instrument to assess functional capacity in dementia: The Texas Functional Living Scale. *Neuropsychiatry, neuropsychology, and behavioral neurology*. 2001. 14: 103-108.
31. Rosca EC, Cornea A, Simu M. Montreal Cognitive Assessment for evaluating the cognitive impairment in patients with schizophrenia: A systematic review. *General Hospital Psychiatry*. 2020. 65: 64-73.
32. Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*. 2005 53: 695-699.

33. Costa AS, Fimm B, Friesen P, Soundjock H, Rottschy C, et al. Alternate-form reliability of the Montreal cognitive assessment screening test in a clinical setting. *Dementia and geriatric cognitive disorders*. 2012. 33: 379-384.
34. Gierus J, Mosiołek A, Koweszko T, Wnukiewicz P, Kozyra O, et al. The Montreal Cognitive Assessment as a preliminary assessment tool in general psychiatry: Validity of MoCA in psychiatric patients. *General hospital psychiatry*. 2015. 37: 476-480.
35. Gil-Berrozpe GJ, Sánchez-Torres AM, García de Jalón E, Moreno-Izco L, Fañanás L, et al. Utility of the MoCA for cognitive impairment screening in long-term psychosis patients. *Schizophrenia Research*. 2020. 216: 429-434.
36. Sieg E, Mai Q, Mosti C, Brook M. The utility of neuropsychological consultation in identifying medical inpatients with suspected cognitive impairment at risk for greater hospital utilization. *The Clinical neuropsychologist*. 2019. 33: 75-89.
37. Adamis D, Helmi L, Fitzpatrick O, Meagher D, McCarthy G. Agreement and equation between Mini Mental State Examination (MMSE) and Montreal Cognitive Assessment (MoCA) in an old age psychiatry outpatient clinic population. *European Psychiatry*. 2016. 33: 184-184.
38. Akbar NL, Effendy E, Camellia V. The Indonesian Version of Montreal Cognitive Assessment (MoCA-Ina): The Difference Scores Between Male Schizophrenia Prescribed by Risperidone and Adjunctive of Donepezil in Public Hospital of Dr Pirngadi Medan, Indonesia. *Open Access Macedonian Journal of Medical Sciences*. 2019. 7: 1762-1767.
39. American Psychiatric Association A. Diagnostic and statistical manual of mental disorders. Washington, DC: American psychiatric association. 1980. 3: 399-406.
40. Arunpongpaissal S, Sangsirilak A. Using MoCA-Thai to evaluate cognitive impairment in patients with schizophrenia. *J Med Assoc Thai*. 2013. 96: 860-865.
41. Barekatin M, Maracy MR, Hassannejad R, Hosseini R. Factors associated with readmission of patients at a university hospital psychiatric ward in Iran. *Psychiatry J*. 2013. 2013: 685625.
42. Bell MD, Bryson G. Work rehabilitation in schizophrenia: does cognitive impairment limit improvement? *Schizophrenia bulletin*. 2001. 27: 269-279.
43. Bergly HT, Sømhovd JM. The Relation Between ADHD Medication and Mild Cognitive Impairment, as Assessed by the Montreal Cognitive Assessment (MoCA), in Patients Entering Substance Use Disorder Inpatient Treatment. *Journal of dual diagnosis*. 14: 228-236.
44. Bezdicek O, Majerova V, Novak M, Nikolai T, Ruzicka E, et al. Validity of the Montreal Cognitive Assessment in the detection of cognitive dysfunction in Huntington's disease. *Applied Neuropsychology: Adult*. 2013. 20: 33-40.
45. Bhugra D. The Global Prevalence of Schizophrenia. *PLoS Med*. 2005. 2: e151.
46. Boone WJ. Rasch Analysis for Instrument Development: Why, When, and How? *CBE Life Sciences Education*. 15.
47. Carson N, Leach L, Murphy KJ. A re-examination of Montreal Cognitive Assessment (MoCA) cutoff scores. *International journal of geriatric psychiatry*. 2018. 33: 379-388.
48. Cruess DG, Sullivan MC, Strainge L, Blackmon JE, Laumann L, et al. Personality predictors of 6-month readmission in adult psychiatric inpatients. *Int J Psychol*. 2022. 57: 613-620.
49. Damian AM, Jacobson SA, Hentz JG, Belden CM, Shill HA, et al. The Montreal Cognitive Assessment and the mini-mental state examination as screening instruments for cognitive impairment: item analyses and threshold scores. *Dementia and geriatric cognitive disorders*. 2011. 31: 126-131.
50. Donisi V, Tedeschi F, Wahlbeck K. Pre-discharge factors predicting readmissions of psychiatric patients: a systematic review of the literature. *BMC Psychiatry*. 16: 449.
51. Feigon S, Hays JR. Prediction of Readmission of Psychiatric Inpatients. *Psychological Reports*. 93: 816-818.
52. Freitas S, Simões MR, Alves L, Santana I. Montreal cognitive assessment: validation study for mild cognitive impairment and Alzheimer disease. *Alzheimer Disease & Associated Disorders*. 27: 37-43.
53. Gerke L, Meyrose K, Ladwig I, Rief W, Nestoriuc Y. Frequencies and Predictors of Negative Effects in Routine Inpatient and Outpatient Psychotherapy: Two Observational Studies. *Frontiers in Psychology*. 2019. 11: 2144.
54. Gold JM, Carpenter C, Randolph C, Goldberg TE, Weinberger DR. Auditory working memory and Wisconsin Card Sorting Test performance in schizophrenia. *Archives of general psychiatry*. 1997. 54: 159-165.
55. Ko RE, Kang D, Park H. Association between the presence of delirium during intensive care unit admission and cognitive impairment or psychiatric problems: the Korean ICU National Data Study. *J intensive care*. 2022. 10: 7.
56. Koski L, Xie H, Finch L. Measuring cognition in a geriatric outpatient clinic: Rasch analysis of the Montreal Cognitive Assessment. *Journal of geriatric psychiatry and neurology*. 2009. 22: 151-160.
57. Lees R, Selvarajah J, Fenton C, Pendlebury ST, Langhorne P, et al. Test accuracy of cognitive screening tests for diagnosis of dementia and multidomain cognitive impairment in stroke. *Stroke*. 2014. 45: 3008-3018.
58. Ma Y, Ji J, Li G, Yang S, Pan S. Effects of donepezil on cognitive functioning the expression level of  $\beta$ -amyloid in peripheral blood of patients with Alzheimer's disease. *Experimental and Therapeutic Medicine*. 15: 1875-1878.
59. Mohamed S, Rosenheck R, Swartz M, Stroup S, Lieberman JA, et al. Relationship of cognition and psychopathology to functional impairment in schizophrenia. *American Journal of Psychiatry*. 2008. 165: 978-987.
60. Murri MB, Folesani F, Costa S, Biancosino B, Colla C, et al. Screening for cognitive impairment in non-affective psychoses: A comparison between the SCIP and the MoCA. *Schizophrenia Research*. 2020. 218: 188-194.
61. Musso WM, Cohen AS, Auster TL, McGovern JE. Investigation of the Montreal Cognitive Assessment (MoCA) as a cognitive screener in severe mental illness. *Psychiatry Research*. 2014. 220: 664-668.
62. Rosca EC, Albarqouni L, Simu M. Montreal cognitive assessment (MoCA) for HIV-associated neurocognitive disorders. *Neuropsychology Review*. 2019. 29: 313-327.
63. Rossetti HC, Lacritz LH, Cullum CM, Weiner MF. Normative data for the Montreal Cognitive Assessment (MoCA) in a population-based sample. *Neurology*. 77: 1272-1275.

64. Rylander M, Colon-Sanchez D, Keniston A, Hamalian G, Lozano A, et al. Risk Factors for Readmission on an Adult Inpatient Psychiatric Unit. *Quality management in health care*. 2016. 25: 22-31.
65. Serper DR, Beech PD, Dill H, Dill C. Neuropsychological and symptom predictors of aggression on the psychiatric inpatient service, *Journal of Clinical and Experimental Neuropsychology*. 2008. 30: 700-709.
66. S, Davidson M. The cost of cognitive impairment in schizophrenia. *Schizophrenia research*. 1995.
67. Smith TE, Hull JW, Romanelli S, Fertuck E, Weiss KA. Symptoms and neurocognition as rate limiters in skills training for psychotic patients. *The American journal of psychiatry*. 156: 1817-1818.
68. Tone HB, Sømhøvd MB. The Relation Between ADHD Medication and Mild Cognitive Impairment, as Assessed by the Montreal Cognitive Assessment (MoCA), in Patients Entering Substance Use Disorder Inpatient Treatment, *Journal of Dual Diagnosis*. 2018. 14: 228-236.
69. US Department of Veteran Affairs, Department of Defense. VA/DoD Clinical Practice Guideline for the Assessment and Management of Patients at Risk for Suicide. 2019.
70. Wood L, Williams C, Enache G, Withers F, Fullarton K, et al. Examining cognitive functioning of adult acute psychiatric inpatients through a brief screening assessment. *Psychiatric Rehabilitation Journal*. 2019. 42: 64-70.
71. Wu C, Dagg P, Molgat C. Measuring stability of cognitive impairment in inpatients with schizophrenia with alternate forms of the Montreal Cognitive Assessment during acute hospitalization. *Psychiatry Research*. 2017. 258: 299-304.
72. Yang Z, Rashid NAA, Quek YF, Lam M, See YM, et al. Montreal cognitive assessment as a screening instrument for cognitive impairments in schizophrenia. *Schizophrenia research*. 2018. 199: 58-63.
73. Zakharova-Luneva E, Cooke DM, Okano S, Hurst C, Geffen S, et al. The relationship between cognition and functional outcomes in rehabilitation: FIM Cog vs. MoCA. *Geriatr Gerontol Int*. 2020. 20: 336-342.
74. Zhang J, Harvey C, Andrew C. Factors Associated with Length of Stay and the Risk of Readmission in an Acute Psychiatric Inpatient Facility: A Retrospective Study. *Australian & New Zealand Journal of Psychiatry*. 2011. 45: 578.